arXiv:1509.01308v2 [math.GR] 24 Sep 2015

Orientable and non-orientable genus n Wicks forms over hyperbolic groups

Andrew Duncan, Steven Fulthorp
School of Mathematics and Statistics
Herschel Building
University of Newcastle upon Tyne
Newcastle upon Tyne

September 25, 2015

[1]

Abstract

In 1962 M.J. Wicks [29] gave a precise description of the form a commutator could take in a free group or a free product and in 1973 extended this description to cover a product of two squares [30]. Subsequently, lists of "Wicks forms" were found for arbitrary products of commutators and squares in free groups and free products [7], [28]. Here we construct Wicks forms for products of commutators and squares in a hyperbolic group. As applications we give explicit lists of forms for a commutator and for a square, and find bounds on the lengths of conjugating elements required to express a quadratic tuple of elements of a hyperbolic group as a Wicks form.

1 Introduction

In 1962 M.J. Wicks [29] described algorithms to decide, given an element u of a free group or a free product of groups, whether or not u is a commutator; and in 1973 [30] extended the results to cover the case where u is a product of two squares. These algorithms are based on the construction of a set of "forms" for commutators and squares. For example, an element u in the free group $F(X)$ is shown to be a commutator if and only if it is conjugate to a cyclically reduced word of the form $F = ABCA^{-1}B^{-1}C^{-1}$, with $A, B, C \in F(X)$ (where cyclically reduced means the word and all its cyclic permutations are reduced). The analogous result for a word u equal to a product a^2b^2 is that u is conjugate to a cyclically reduced word $F_1 = A^2BC^2B^{-1}$ or $F_2 = ABACB^{-1}C$. We call the words F, F_1 and F_2 *Wicks forms* for commutators and products of two squares, in free groups. The lists of forms [29], [30] for commutators and products of two squares in free products of groups are similar though more involved.

---

[1] AMS Mathematics Subject Classification: 20F12, 20F65, 20F67

The question of whether an element is a commutator or product of squares is an example of an equation over a group: as defined in Section 2.3. Landmarks in the theory of equations over groups are the papers of Makanin [23] showing that arbitrary systems of equations over a free group are solvable, in the free group; and of Razborov [26] giving a parametric description of the set of all solutions. The complexity of Makanin's algorithm is very high: it is not known to be primitive recursive, whereas Wicks' algorithms which show that the quadratic equations $[a, b] = u$ and $a^2b^2 = u$, in variables a, b, are solvable in free groups and free products, run in time bounded by a polynomial in the length of u. In addition, quadratic equations play an important part in algorithms for solution of general systems of equations, and they are closely related to compact surfaces. For these reasons, among others, the study of quadratic equations is of independent interest.

An element of a group is said to have *orientable genus* n if it may be expressed as a product of n commutators, and no fewer, and *non-orientable genus* $n/2$ if it is a product of n squares, and no fewer (see Section 2.3 below). The corresponding quadratic equations were shown to be solvable in a free groups by Edmunds [10], [11]. Culler [8] and, independently, Goldstein and Turner [13], used the theory of compact surfaces to produce such algorithms; in particular Culler described a set of Wicks forms for elements of orientable and non-orientable genus n in the free group. Algorithms to decide the solvability, and to find a solution, of a general quadratic equation over a free group and over a free product of groups (where quadratic equations are solvable in the factors) were given by Comerford and Edmunds [5]. In the case of the free group, Ol'shanski [24] (see also [15]) shows that there is such an algorithm which runs in time polynomial in the length of the coefficients of the equation (see Section 2.3 for more detail). The set of all solutions of a quadratic equation in a free group $F(X)$ can, as shown by Comerford and Edmunds [6], be described in terms of a finite set of *basic* solutions and certain $F(X)$-automorphisms of $F(\mathcal{A}) * F(X)$; and all the parameters of this description may be effectively computed, given the equation $w = 1$. In fact Grigorchuk and Kurchanov [15] show that the set of basic solutions of the quadratic equation $w = 1$ may be computed in polynomial time in the length of the coefficients of w. Moreover in [15] a set of Wicks forms for solutions of a given quadratic equation in a free group is shown to exist.

Quadratic equations in small cancellation groups and hyperbolic groups were studied in [25], [18], [22]. Ol'shanskii [24][Theorem 6] (see also [15]) constructed an algorithm, to determine whether or not a quadratic equation of genus g has solution in H, and if so to find one. Moreover this algorithm runs in time bounded by a polynomial in the sum of the lengths of coefficients. A parametric description of the set of solutions of a quadratic equation was described in [14]. Generalising Makanin's algorithm, Rips and Sela [27] proved the existence of an algorithm for the solvability of a finite system of equations over a torsion-free hyperbolic group, and Dahmani and Guirardel [9] have extended this result to cover the case of hyperbolic groups with torsion.

In [28] Vdovina described a procedure for constructing Wicks forms for elements of any orientable genus in a free product. In this paper we use similar methods to construct Wicks forms for elements of arbitrary genus in hyperbolic groups. Using the forms of genus 1 and 1/2 we then list all the possible forms of commutators and squares in a hyperbolic group. Similar lists of forms could be constructed for elements of higher genus. However, the number of possible extensions increases dramatically with the increase of genus: Bacher and Vdovina show [2] that the number of orientable Wicks forms of genus g is $\Omega(g!)$. We also describe, as an application of our results, forms for quadratic tuples of words (defined below).

We begin by introducing a number of definitions and preliminary results in Sections 2.1 to 2.3. The key technique which we call an "extension" of a quadratic word is introduced and developed in Sections 2.5 and 2.6. This puts us in a position to state the main results of the paper: Theorem 2.23 and Theorem 2.24, in Section 2.7. Section 3 contains some applications of these theorems. In Sections 3.1 and 3.2, we prove Propositions 3.1 and Propositions 3.2, which describe the possible forms for commutators and squares in a δ-hyperbolic group H. In Section 3.3 we discuss quadratic equations, give bounds on the length of elements in "minimal" solutions to such equations and show how the main theorems may be used to describe forms for quadratic tuples of elements. The proofs of the main theorems are left to Section 4. In Section 4.1 we give preliminary results needed to construct an appropriate extension; and complete this construction in Section 4.2. Finally, in Section 4.4 we bound the length of the conjugator which appears in the main theorems.

2 Definitions and Main results

2.1 Hyperbolic groups

For background on hyperbolic groups the reader is referred to [1, 12, 17]. Here we use the notation of [1]. The Cayley graph of a group G with respect to a generating set X will be denoted $\Gamma_X(G)$. As usual, the word metric and makes G into a geodesic metric space. We use $[x, y]$ to denote a geodesic path p from x to y and $(x, y]$, $[x, y)$ and (x, y) to denote $p\backslash\{x\}$, $p\backslash\{x\}$ and $p\backslash\{x, y\}$, respectively. If p is a path from x to y then we write $x = \iota(p)$, $y = \tau(p)$, we denote the reverse of p from y to x by p^{-1} and, if p is geodesic, then we write $|p| = d(x, y)$. Let w be a (reduced) word in $F(X)$. The length of w as a word is denoted $|w|$. If $|v| \geq |w|$ for all words v in $F(X)$ such that $w =_G v$ then we say that the word w is *G-minimal*. We use the notation $|w|_G$ to denote the length of a G-minimal word in $F(X)$ which is equal to w in G. It follows that, for all $g \in G$, a G-minimal word representing g is the label of a geodesic path from 1 to g in $\Gamma_X(G)$.

A *geodesic triangle*, denoted $\triangle xyz$, consists of the union of three points x, y, z, of the the Cayley graph, and geodesic paths $[x, y]$, $[y, z]$ and $[x, z]$ joining them. Let H be a group generated by a set X and let $T = \triangle xyz$ be a geodesic triangle in $\Gamma_X(H)$. Let $T' = \triangle' x'y'z'$ be a Euclidean triangle with sides the same length as T. That is $d_E(x', y') = d(x, y)$ etc, where d_E is the standard Euclidean metric. There is a natural identification map ϕ from T to T'. The maximum inscribed circle in T' meets the side $[x'y']$ (respectively $[y'z']$, $[z'x']$)

at a point c_z (respectively c_x, c_y) such that

$$d(x', c_z) = d(x', c_y), d(y', c_x) = d(y', c_z), d(z', c_y) = d(z', c_x).$$

Notice that we have

$$d(x', c_z) = \frac{1}{2}(d(x', c_z) + d(x', c_y)) = \frac{1}{2}(d(x', z') + d(x', y') - d(z', y')). \tag{2.1}$$

The preimages $\phi^{-1}(c_x)$, $\phi^{-1}(c_y)$ and $\phi^{-1}(c_z)$ of c_x, c_y and c_z in T are called the *internal points* of T.

There is a unique isometry ι_T of the triangle T' onto a *tripod*: that is a tree with one vertex p of degree three and vertices x'', y'', z'' each of degree one, such that $d(p, z'') = d(z', c_y) = d(z', c_x)$ etcetera. Let ψ be the composite map $\psi = \iota_T \circ \phi$. We say that a geodesic triangle is δ*-thin* if the fibres of ψ have diameter at most δ in $\Gamma_X(H)$. That is, for all x, y in T,

$$\psi(x) = \psi(y) \implies d(x, y) \le \delta. \tag{2.2}$$

Definition 2.1. *A group H is said to be* δ-hyperbolic, *with respect to a finite generating set X, if all geodesic triangles in $\Gamma_X(H)$ are δ-thin. A group is* (word) hyperbolic *if, for some $\delta \ge 0$, it is δ-hyperbolic, with respect to some generating set X.*

In the light of (2.1) and (2.2) a group is δ-hyperbolic if and only if it satisfies the following condition. Let w and z be any words in $F(X)$ which are H-minimal, with $w = w_1 w_2$ and $z = z_1 z_2$, $|w| = |w_1| + |w_2|$ and $|z| = |z_1| + |z_2|$.

$$\text{If } |w_2| = |z_1| \le \frac{1}{2}(|w| + |z| - |wz|_H) \text{ then } |w_2 z_1|_H \le \delta. \tag{2.3}$$

It can be shown that word hyperbolic groups are finitely presented. For this, and an account of many characterisations of hyperbolic groups, see, for example, [1]. For the remainder of the paper, H will denote a δ-hyperbolic group, with respect to the generating set X, and we shall assume that H has a finite presentation $\langle X|S\rangle$.

The following lemma of Gromov (see [14]) shows that the conjugacy problem is solvable in hyperbolic groups.

Lemma 2.2 ([14])**.** *If H-minimal words h_1 and h_2 are conjugate in H, then a word w can be found such that $h_1 =_H w h_2 w^{-1}$ and*

$$|w| \le \frac{1}{2}(|h_1| + |h_2|) + M + 1,$$

where M is the number of elements of H represented by words of $F(X)$ of length $\le 4\delta$.

(The obvious algorithm for the conjugacy problem based on the lemma above takes exponential time, in the length of the two input words, but a closer analysis yields linear bounds on the time required: see Bridson and Haefliger [4][pp. 451–454] or Bridson and Howie [3].)

2.2 Thin polygons

Let $H = \langle X|S\rangle$ be a finitely generated hyperbolic group. The following lemma is a more explicit version of [16][6.1C]; and of [24][Lemma 10]. Its statement requires some further notation. Let $\gamma = [x, y]$ be a geodesic in $\Gamma_X(H)$ and let $v_0, \dots, v_n$ be points of γ such that $v_0 = x$, $d(x, v_{i-1}) < d(x, v_i)$ and $v_n = y$, and set $\gamma_i = [v_{i-1}, v_i]$, $i = 1, \dots, n$. Then the sequence $\gamma_1, \dots, \gamma_n$ is called a *partition* of γ. If $\gamma = [x, y]$ and $\gamma' = [x', y']$ are geodesics such that $|\gamma| = |\gamma'|$ and, for all $z \in \gamma$ and $z' \in \gamma'$, if $d(x, z) = d(x', z')$ then $d(z, z') \le k$; we say that γ and γ' *k-fellow travel.*

Lemma 2.3. *Let $q = \gamma_0\gamma_1 \dots \gamma_n$ be a closed path in $\Gamma_X(H)$, where γ_i is a geodesic path, for $i = 0, \dots, n$, $n \ge 1$. Then*

- *there exist integers $r_0, \dots, r_n$, $2 \le r_i \le n$ and, for $i = 0, \dots n$, there exists a partition $\gamma_i^{(1)}, \dots, \gamma_i^{(r_i)}$ of γ_i and,*
- *setting $I = \cup_{i=0}^n \{(i, j) : 1 \le j \le r_i\}$, there exists an involution σ of I,*

such that the following hold.

(i) Writing $\sigma(0, i) = (a_i, b_i)$, for $1 \le i \le r_0$, geodesics $\gamma_0^{(i)}$ and $(\gamma_{a_i}^{(b_i)})^{-1}$ are $\delta\lceil\log_2(n)\rceil$-fellow travellers and $n \ge a_1 \ge \dots \ge a_{r_0} \ge 1$; and

(ii) fixing i with $1 \le i \le n$ and writing $\sigma(i, j) = (a_j, b_j)$, geodesics $\gamma_i^{(j)}$ and $(\gamma_{a_j}^{(b_j)})^{-1}$ are $\delta(2\lceil\log_2(n)\rceil - 1)$-fellow travellers and there exists an integer c_i, such that $1 \le c_i \le r_i - 1$ and

$$n \ge a_{c_i+1} \ge \cdots \ge a_{r_i} \ge i + 1 \text{ and } i - 1 \ge a_1 \ge \cdots \ge a_{c_i} \ge 0.$$

(iii) In addition, $\sigma(0, 1) = (n, n)$, $\sigma(i, 1) = (i - 1, n)$, for $1 \le i \le n$, and $\gamma_i^{(1)}$ and $(\gamma_{i-1}^{(r_{i-1})})^{-1}$ are δ-fellow travellers, for $0 \le i \le n$ (subscripts modulo $n + 1$).

(Here $\iota(\gamma_i^{(j)})$ and $\tau(\gamma_i^{(j)})$ are not necessarily vertices of $\Gamma_H(X)$. In (ii) if $a_j = 0$ then, from (i), in fact $\gamma_i^{(j)}$ and $(\gamma_{a_j}^{(b_j))^{-1}}$ are $\delta\lceil\log_2(n)\rceil$-fellow travellers.)

An instance of the case $n = 4$ is illustrated in Figure 2.1.

Proof. Let $k = \lceil\log_2(n)\rceil$. By adding paths of length zero between $\tau(\gamma_n)$ and $\iota(\gamma_0)$, we may assume that $n = 2^k$. Thus we may replace $\lceil\log_2(n)\rceil$ with k, throughout the statement of the lemma. If $k = 1$ then the result follows directly from the thin triangles condition: the internal points of a triangle determine appropriate partitions of the sides.

For the induction we need to be able to modify the partitions of the γ_i by adding new points, so we begin by describing how to do this. The data consisting of the partitions $\gamma_i^{(1)}, \dots, \gamma_i^{(n)}$ of the γ_i and the map σ, as in the lemma, is called a *matching* of q, *based* at γ_0. Given a matching M as described in the statement of the lemma, satisfying (i), (ii) and (iii), supp pose that for some i, j we have points p, r such that $\gamma_i^{(j)} = [p, r]$ and a point



Figure 2.1: $\gamma_2^{(3)}$ and $(\gamma_4^{(2)})^{-1}$ are 4δ-fellow travellers and $\gamma_0^{(2)}$ and $(\gamma_2^{(2)})^{-1}$ are 2δ-fellow travellers. All other paired intervals δ-fellow travel.

$q \in [p, r]$, $p \neq q \neq r$. We form a *simple refinement* M' of M as follows. Let $\sigma(i, j) = (a, b)$, let $\gamma_a^{(b)} = [p', r']$ and let q' be the point of $[p', r']$ such that $d(p, q) = d(q', r')$. Replace the partition $\gamma_i^{(1)}, \ldots, \gamma_i^{(r_i)}$ of γ_i with the partition $\gamma_i'^{(1)}, \ldots, \gamma_i'^{(r_i+1)}$, where $\gamma_i'^{(s)} = \gamma_i^{(s)}$, for $1 \leq s \leq j-1$, $\gamma_i'^{(j)} = [p, q]$, $\gamma_i'^{(j+1)} = [q, r]$ and $\gamma_i'^{(s)} = \gamma_i^{(s-1)}$, for $j+2 \leq s \leq r_i+1$. Replace the partition $\gamma_a^{(1)}, \ldots, \gamma_a^{(r_a)}$ of γ_a with the partition $\gamma_a'^{(1)}, \ldots, \gamma_a'^{(r_a+1)}$, formed in exactly the same way (replacing i, j, p, q and r, by a, b, p', q' and r', throughout). Now replace σ with an involution σ' such that

$$\begin{array}{ll}
\sigma'(i,s) = \sigma(i,s), 1 \leq s \leq j-1, & \sigma'(a,t) = \sigma(a,t), 1 \leq t \leq b-1 \\
\sigma'(i,j) = (a, b+1) \text{ and } \sigma'(i,j+1) = (a,b), & \sigma'(a,b) = (i, j+1) \text{ and } \sigma'(a,b+1) = (i,j) \\
\sigma'(i,s) = \sigma(i,s-1), j+2 \leq s \leq r_i+1, & \sigma'(a,t) = \sigma(a,t-1), b+2 \leq t \leq r_b+1,
\end{array}$$

and for $u \neq i, a$,

$$\sigma'(u,v) = \begin{cases} (i, s+1) & \text{if } \sigma(i,s) = (u,v), \text{ where } j+1 \leq s \leq r_i, \\ (a, t+1) & \text{if } \sigma(a,t) = (u,v), \text{ where } b+1 \leq t \leq r_a, \\ \sigma(u,v) & \text{otherwise.} \end{cases}$$

Then these partitions of the γ_i together with the function σ' form a new matching of q, which again satisfies conditions (i), (ii) and (iii). A finite sequence of simple refinements of M is called a *refinement*.

Assume then that $k > 1$, that the result holds for all non-negative integers n no larger than 2^{k-1} and that $n = 2^k$. Let γ_0' be the geodesic path from $\tau(\gamma_{2^{k-1}})$ to $\tau(\gamma_0)$ and let $\gamma_{2^{k-1}}'$ be the geodesic path from $\iota(\gamma_0)$ to $\tau(\gamma_{2^{k-1}})$ (see Figure 2.2). Let c_0, c_0' and c_0'' be the internal points of triangle $T = \gamma_0 \gamma_0'^{-1} \gamma_{2^{k-1}}'^{-1}$, with c_0 on γ_0, c_0' on γ_0' and c_0'' on $\gamma_{2^{k-1}}'$ and let $t_0 = \iota(\gamma_0)$, $t_1 = \tau(\gamma_0)$ and $t_2 = \iota(\gamma_0')$ be the vertices of T. Let M_0 be the corresponding



Figure 2.2

matching, with partitions $[t_0, c_0], [c_0, t_1]$ of γ_0; $[t_1, c'_0], [c'_0, t_2]$ of γ'^{-1}_0 and $[t_2, c''_0], [c''_0, t_0]$ of $\gamma'^{-1}_{2^{k-1}}$; and involution σ_0 such that $\sigma_0(0,1) = (2,2)$, $\sigma_0(0,2) = (1,1)$ and $\sigma_0(1,2) = (2,1)$. Let $q'_1 = \gamma'_0\gamma_1 \cdots \gamma_{2^{k-1}}$ and $q'_2 = \gamma'_{2^{k-1}}\gamma_{2^{k-1}+1} \cdots \gamma_{2^k}$. From the inductive hypothesis, there exist matchings M_1 of q'_1 and M_2 of q'_2, based at γ'_0 and $\gamma'_{2^{k-1}}$, respectively. Suppose that σ_i is the involution associated to M_i and that, under these matchings the partitions of γ'_0 and $\gamma'_{2^{k-1}}$ are $\gamma'^{(1)}_0, \dots, \gamma'^{(r')}_0$ and $\gamma'^{(1)}_{2^{k-1}}, \dots, \gamma'^{(r'')}_{2^{k-1}}$, respectively.

We shall refine the partitions M_1 and M_2, so that the intervals, lying between t_2 and either c'_0 or c''_0, are of the same number and lengths. First, single out the end points of intervals of the partitions of γ'_0 and $\gamma'_{2^{k-1}}$ which lie between $[t_2, c'_0]$ and $[c''_0, t_2]$. That is, let $S_1 = \{p \in [t_2, c'_0] : p = \tau(\gamma'^{(j)}_0), 1 \le j \le r'\}$ and let $S_2 = \{p \in [c''_0, t_2] : p = \iota(\gamma'^{(j)}_{2^{k-1}}), 1 \le j \le r''\}$. Now form sets of points to be used in the refinement: let $T_1 = \{q \in [t_2, c'_0] : d(q, t_2) = d(p, t_2),$ for some $p \in S_2\}$ and $T_2 = \{q \in [c''_0, t_2] : d(q, t_2) = d(p, t_2),$ for some $p \in S_1\}$. Refine M_1 by adding the points of $T_1 \cup \{c'_0\}$ to γ'_0 and refine M_2 by adding the points of $T_2 \cup \{c''_0\}$ to $\gamma'_{2^{k-1}}$. We shall now consider M_1 and M_2 to be these refined matchings and, after adjusting notation, take the partitions of γ'_0 and $\gamma'_{2^{k-1}}$ to be $\gamma'^{(1)}_0, \dots, \gamma'^{(r')}_0$ and $\gamma'^{(1)}_{2^{k-1}}, \dots, \gamma'^{(r'')}_{2^{k-1}}$, again.

Now we refine M_0, using the partitions of γ'_0 and $\gamma'_{2^{k-1}}$ under M_1 and M_2. Let $U_1 = \{p \in [t_2, t_1] : p = \tau(\gamma'^{(j)}_0), 1 \le j \le r'\}$ and $U_2 = \{p \in [t_0, t_2] : p = \iota(\gamma'^{(j)}_{2^{k-1}}), 1 \le j \le r''\}$. Refine M_0 by adding the points of $U_1 \cup U_2$. As before we refer to the refinement as M_0 and to its involution as σ_0. Under M_0 we have a partition $\gamma^{(1)}_0, \dots, \gamma^{(r_0)}_0$ of γ_0 with the following properties. For j such that $\tau(\gamma^{(j)}_0) \in [t_0, c_0]$ the interval $\gamma^{(j)}_{2^{k-1}}$ has terminal point on $[t_0, c''_0]$ and δ-fellow travels with $\gamma^{(j)}_0$. From the inductive hypothesis, and the fact that refinements preserve properties (i), (ii) and (iii), $\gamma^{(j)}_{2^{k-1}}$ and $(\gamma^{(b)}_a)^{-1}$ are $\delta(k-1)$-fellow travellers, where $(a, b) = \sigma_2(2^{k-1}, j)$. Hence $\gamma^{(j)}_0$ and $(\gamma^{(b)}_a)^{-1}$ are δk-fellow travellers and we may set $\sigma(0, j) = \sigma_2(2^{k-1}, j)$. Similarly, for j such that $\iota(\gamma^{(j)}_0) \in [c_0, t_1]$, if $\sigma_1(0, r' - r_0 + j) = (a, b)$ then the intervals $\gamma^{(j)}_0$ and $(\gamma^{(b)}_a)^{-1}$ are δk-fellow travellers and we may set $\sigma(0, j) = \sigma_1(0, r' - r_0 + j)$. This defines σ on all pairs $(0, j)$, with $1 \le j \le r_0$. If $1 \le i \le 2^{k-1}$ and j is such that

$(i,j) \neq \sigma(0,j')$, for all j', it follows from construction of M_1 that $\sigma_1(i,j) = (a,b)$ where either $a = 0$ and ${\gamma'}^{(b)}_0$ has initial point in $[t_2, c'_0]$ or $1 \leq a \leq 2^{k-1}$. In the former case ${\gamma'}^{(b)}_0$ is a subinterval of $[t_2, c'_0]$, which $\delta(k-1)$-fellow travels with $(\gamma^{(j)}_i)^{-1}$, $\sigma_0(0,b) = (2^{k-1}, l)$, and the interval ${\gamma'}^{(l)}_{2^{k-1}}$ of $\gamma'_{2^{k-1}}$ is a subinterval of $[c''_0, t_2]$ and δ-fellow travels with $({\gamma'}^{(b)}_0)^{-1}$. Furthermore, $\sigma_2(2^{k-1}, l) = (c,d)$, for some c with $2^{k-1}+1 \leq c \leq 2^k$, and ${\gamma'}^{(l)}_{2^{k-1}}$ and $(\gamma^{(d)}_c)^{-1}$ are $\delta(k-1)$-fellow travellers. Hence, in this case, $\gamma^{(j)}_i$ and $(\gamma^{(d)}_c)^{-1}$ are $2\delta(k-1)$-fellow travellers; and we may set $\sigma(i,j) = (c,d)$. In the latter case $\sigma_1(i,j) = (a,b)$, where $1 \leq a \leq 2^{k-1}$, so $\gamma^{(j)}_i$ and $(\gamma^{(b)}_a)^{-1}$ are $\delta(2(k-1)-1)$-fellow travellers, so we may set $\sigma(i,j) = (a,b)$. This defines σ on all pairs (i,j), where $1 \leq i \leq 2^{k-1}$. A similar argument allows us to define σ on the remaining pairs (i,j) where $2^{k-1}+1 \leq i \leq 2^k$. Therefore we have a matching M which satisfies (i) and (ii). If this matching does not already satisfy (iii) then a simple modification results in one that does. □

2.3 Quadratic equations, genus and quadratic words

We write $\mathbb{Z}[\frac{1}{2}]$ for the set $\{n/2 : n \in \mathbb{Z}\}$. We use Stallings' definition of the genus of a surface: a connected sum of g torii has *genus* g, while a connected sum of g projective planes has *genus* $g/2$. (Hence the Euler characteristic of a compact, connected, closed surface Δ, of genus g, is $\chi(\Delta) = 2 - 2g$, irrespective of orientability.)

By a *graph* we mean a finite directed graph. If e is an edge of a graph we write $\iota(e)$ and $\tau(e)$ for the initial and terminal vertices of e. We use e^{-1} to denote the edge with $\iota(e^{-1}) = \tau(e)$ and $\tau(e^{-1}) = \iota(e)$. If p is a path in a graph with edge sequence $e_1, \dots, e_n$ then we denote by p^{-1} the path with edge sequence $e_n^{-1}, \dots, e_1^{-1}$. A *labelling* of a graph is a function λ from directed edges of the graph to a set L such that if e is an edge with $\lambda(e) = A$ then $A^{-1} \in L$ and $\lambda(e^{-1}) = A^{-1}$. We routinely identify the edge e with its label $\lambda(e)$.

Let $\mathcal{A}$ be an countably infinite alphabet. The *support* of a word w in the free monoid $(\mathcal{A} \cup \mathcal{A}^{-1})^*$ is the smallest subset $\mathcal{S}$ of $\mathcal{A}$ such that w belongs to $(\mathcal{S} \cup \mathcal{S}^{-1})^*$, written $\operatorname{supp}(w)$. As we are concerned here only with groups, we refer to elements of $(\mathcal{A} \cup \mathcal{A}^{-1})^*$ as *words over* $\mathcal{A}$. A word in $(\mathcal{A} \cup \mathcal{A}^{-1})^*$ is *(freely) reduced* if it contains no subword $x^{\varepsilon}x^{-\varepsilon}$, where $x \in \mathcal{A} \cup \mathcal{A}^{-1}$, $\varepsilon = \pm 1$, and *(freely) cyclically reduced* if w and all its cyclic permutations are reduced. We use the term *cyclic word* to mean the equivalence class $[w]$ of a word w under the relation which relates two words if one is a cyclic permutation of the other. When we talk about a *cyclic word* w we mean any element v of $[w]$.

Definition 2.4. *Let $w_1, \dots, w_t$ be a t-tuple of words over $\mathcal{A}$ and let $W = w_1 \cdots w_t$ (as a word in $(\mathcal{A} \cup \mathcal{A}^{-1})^*$). Then $w_1, \dots, w_t$ is said to be* quadratic *if each element of* $\operatorname{supp}(W)$ *appears exactly twice in W. In this case, let $x \in$* $\operatorname{supp}(W)$ *and let $W = W_0 x^{\varepsilon} W_1 x^{\delta} W_2$, where $\varepsilon, \delta \in \{\pm 1\}$. Then the* signature $\sigma(x)$ of x in $w_1, \dots, w_t$ *is $\sigma(x) = (\varepsilon, \delta)$. Define $o(x) = -\varepsilon\delta$, where $\sigma(x) = (\varepsilon, \delta)$. If $o(x) = 1$ then we say that the letter x is* alternating, *in $w_1, \dots, w_t$. If every element x of* $\operatorname{supp}(W)$ *is alternating then $w_1, \dots, w_t$ is said to be* orientable. *The* signature *of a quadratic tuple $w_1, \dots, w_t$ is the map σ from* $\operatorname{supp}(W)$ *to* $\{\pm 1\} \times \{\pm 1\}$.

Let X and $\mathcal{A}$ be disjoint sets and let w be an element of $F(X) * F(\mathcal{A})$. The expression $w = 1$ is called an *equation*, with variables $\mathcal{A}$. If H is a group with presentation $\langle X|S\rangle$ then a homomorphism ϕ from $F(X) * F(\mathcal{A})$ to H is an *H-map* if the restriction of ϕ to $F(X)$ induces the identity map on H. A *solution* to the equation $w = 1$ over H is an H-map $\phi : F(X) * F(\mathcal{A}) \to H$ such that $\phi(w) =_H 1$. Given $w \in F(X) * F(\mathcal{A})$, write $w = c_0a_1c_1 \dots c_{n-1}a_n$, where $a_i \in F(\mathcal{A})$ and $c_i \in F(X)$, with $c_i \neq 1$, $i > 0$, and $a_i \neq 1$, $i < n$. Then the equation $w = 1$ is said to have *coefficients* $c_0, \dots c_{n-1}$. The equation $w = 1$, with $w = c_0a_1c_1 \dots c_{n-1}a_n$ as above, is *quadratic* if $a_1, \dots, a_n$ is a quadratic n-tuple over $\mathcal{A}$, *orientable* if $a_1, \dots, a_n$ is orientable, and *non-orientable* otherwise.

To a quadratic equation $w = 1$, with coefficients $c_0, \dots, c_{n-1}$, we associate a surface Σ as follows. Take a copy D of the disk D^2 and divide its boundary into $|w|$ segments (the length taken in $F(\mathcal{A} \cup X)$). Write w anti-clockwise around the boundary of a disk, labelling segments consecutively with the letters of $\mathrm{supp}(w)$, and directing segments anti-clockwise when a letter occurs with exponent 1, and clockwise when the labelling letter has exponent -1. For example if $w = ABCA^{-1}B^{-1}C^{-1}$, where $A, B, C \in \mathcal{A}$, we obtain the labelling at the top left of Figure 2.4. Now identify the segments labelled by the same letters, respecting orientation. We obtain a compact surface Σ_w of genus g, for some $g \geq 0$, $g \in \mathbb{Z}[\frac{1}{2}]$, with n boundary components, which is orientable if and only if $w = 1$ is orientable. After identification the oriented segments on the boundary of the disk become the edges of a directed, labelled graph Γ_w on the surface Σ_w; with boundary if $n - 1 > 0$. We call this graph the *graph associated to w*. (See Example 2.6.) Every edge e of Γ_w is labelled with an element of $\mathrm{supp}(w)$ and directed edges are in one to one correspondence with $\mathrm{supp}(w)$; so we identify edges with $\mathrm{supp}(w)$. Edges on the boundary of Σ are labelled by elements of X while two-sided edges are labelled by elements of $\mathcal{A}$. The equation $w = 1$ is said to have *genus* g if Σ is a surface of genus g.

Now consider the following types of quadratic equation. The orientable, genus g, quadratic equation

$$\prod_{i=1}^{t} v_i^{-1}c_iv_i \prod_{i=1}^{g}[x_i, y_i] = 1, \tag{2.4}$$

with variables v_i, x_i, y_i, and the non-orientable, genus g, quadratic equation

$$\prod_{i=1}^{t} v_i^{-1}c_iv_i \prod_{i=1}^{2g} z_i^2 = 1, \tag{2.5}$$

with variables v_i, z_i, where the coefficients are the c_i in both cases, are called *standard* quadratic equations. A quadratic equation $w = 1$ is equivalent to a standard quadratic equation: in the sense that there is an efficient procedure to transform $w = 1$ into one of (2.4) or (2.5), and to transform a solution ϕ of $w = 1$ into a solution of the resulting equation; and vice-versa (see for example [14][Proposition 2.2]).

Now let $w = 1$ be the equation (2.4) or (2.5) and suppose ψ is a solution to $w = 1$ over H. In the construction of the surface Σ_w above, we labelled the boundary of a disk D with the word w. Replacing the label a of each directed edge of D with $\psi(a)$ we obtain a disk with boundary label $\psi(w)$. As $\psi(w) =_H 1$, there exists a van Kampen diagram on D, with boundary label $\psi(w)$, over H; and this gives a van Kampen diagram on Σ_w.

Conversely, consider a van Kampen diagram K on a genus g, surface Σ with t boundary components. To construct a solution to one of (2.4) or (2.5) we allow van Kampen diagrams to contain *null-cells*: that is 2-cells with boundary labels of the form $1 \cdot x \cdot 1 \cdot x^{-1}$, $x \in X$. Given a 2-cell R of K we may replace each directed edge of R with boundary label x by a null-cell; as on the left hand side of Figure 2.4. We call the result a *thickening* of R. Whenever we do this it gives us a new van Kampen diagram, on Σ with the same boundary labels. Returning to K, we wish to show that we may cut Σ along a closed curve c which separates Σ into two surfaces, such that capping each surface off with a disk glued along its copy of c, one surface Σ_g is of closed of genus g and the other is a sphere S with t boundary components, labelled $c_1, \dots, c_t$, when read with the appropriate orientation of S. Moreover, we would like to choose c to be a subgraph of the 1-skeleton K_1 of K. Since Σ is the 2-skeleton of K, the graph K_1 is connected, and we may choose a minimal connected subgraph T of K_1 which contains every boundary component. Thicken every 2-cell which meets T. The union of the null-cells of (this thickened version of) K meeting T is then a genus 0 sub-surface Δ of Σ containing all boundary components. The boundary of Δ has no self intersections and is a path in K_1. If Σ is orientable and the boundary components are labelled $c_1, \dots, c_t$, when read with some chosen orientation of Σ then the boundary components of Δ are also labelled $c_1, \dots, c_t$; and so we may take $c = \delta\Delta$.

In the case that Σ is non-orientable and $t > 0$, our assumption on the boundary components means that each boundary component may be assigned an orientation in such a way that, reading each boundary component's label, with its chosen orientation, from a suitable base point b_i, the boundary labels are $c_1, \dots, c_t$. This means in turn that, reading the labels of the boundary components of the disk Δ constructed above, according to some fixed orientation of Δ, we obtain the list $c_1^{\varepsilon_1}, \dots, c_t^{\varepsilon_t}$, with $\varepsilon_i = \pm 1$ (ignoring the boundary label of c itself). To modify this disk so that the ε_i are the same, for all i, we use the non-orientability of Σ. First choose a vertex $*$ of K on the boundary c of the disk D, and a base point b_i on the boundary component labelled c_i, for each i (so that the boundary label may be read from b_i). Then, after thickening some 2-cells if necessary, we may choose paths p_i in $D \cap K_1$, from $*$ to p_i, for $i = 1, \dots, t$, so that the p_i have disjoint interiors and the boundary component of the surface obtained by cutting Σ along the p_i is $\prod_{i=1}^t w_i c_i w_i^{-1}$, where w_i is the label of p_i, as in Figure 2.3a.

Suppose that $\varepsilon_1 = 1$ and i is minimal such that $\varepsilon_i = -1$. Then choose vertices v_1 and v_2 on c and paths q_1, from $*$ to v_1, and q_2, from v_2 to b_i, so that the path $q_1^{-1} p_i q_2^{-1}$ cuts a disk from D, not meeting any boundary component c_j and so that the q_1 meets p_j only at $*$, for all j, and q_2 meets p_i only at b_i and meets no other p_j; as in Figure 2.3b. Now let q be a simple, orientation reversing, path on Σ from v_1 to v_2 such that q meets D only at v_1 and v_2, as in Figure 2.3c. Such a path exists since Σ is non-orientable. After thickening the 2-cells meeting these paths we may assume that q and q_i are edge paths in K_1. Replace the path p_i with $p_i' = q_1 q q_2$. Now repeat the construction of D using the tree $p_1 \cup \cdots \cup p_i' \cup \cdots \cup p_t$ instead of T. This time the boundary components have labels $c_1^{\varepsilon_1}, \dots, c_t^{\varepsilon_t}$, where $\varepsilon_1 = \cdots = \varepsilon_i = 1$. Continuing this way, eventually all the ε_i are equal.



Figure 2.3: Reversing orientation of a boundary component

Thus, in all cases, there is a closed edge path c in K_1 with the required properties. Let C be the label of c. The disk Δ gives a solution to the equation $\prod_{i=1}^{t} v_i^{-1}c_iv_i = C$ and the surface $\Sigma\backslash\mathrm{int}(\Delta)$ gives a solution to the equation $\prod_{i=1}^{g}[x_i, y_i] = C^{-1}$, if Σ is orientable, and $\prod_{i=1}^{2g} z_i^2 = C^{-1}$, otherwise. Combining these we obtain a solution to (2.4) or (2.5) as appropriate. Thus there exists a solution to $w = 1$ if and only if there exists a van Kampen diagram on a surface Σ, of genus g, with t boundary components, labelled (with appropriate orientation) $c_1, \dots, c_t$. This motivates the following definition.

Definition 2.5. *Let G be a group and let $c_1, \dots, c_t$ be elements of G.*

1. *The* orientable genus *of $(c_1, \dots, c_t)$, denoted* $\mathrm{genus}_G^+(c_1, \dots, c_t)$*, is the smallest integer k such that there exists a solution to the equation* (2.4) *over G.*
2. *The* non-orientable genus *of $(c_1, \dots, c_t)$, denoted* $\mathrm{genus}_G^-(c_1, \dots, c_t)$*, is the smallest element $k \in \mathbb{Z}[\frac{1}{2}]$ such that there exists a solution to the equation* (2.4) *over G.*

In all cases, if there is no k satisfying the given conditions then the corresponding genus is defined to be infinite.

When no ambiguity arises, or we wish to make statements covering both orientable and non-orientable genus we use $\mathrm{genus}_G(c_1, \dots, c_t)$ instead of $\mathrm{genus}_G^+(c_1, \dots, c_t)$ or $\mathrm{genus}_G^-(c_1, \dots, c_t)$. (2.4) holds, and similarly for genus_G^-.

2.4 Wicks forms

Consider a quadratic word U over $\mathcal{A}$ and the surface Σ_U obtained as above. The signature σ and orientability o of letters of U induce a signature and orientability on the graph Γ_U: namely if $\lambda(e) = A \in \mathrm{supp}(U)$ then $\sigma(e) = \sigma(A)$ and $o(e) = -\varepsilon\delta$, where $\sigma(A) = (\varepsilon, \delta)$. The edge e is *alternating* if $o(e) = 1$. To obtain a van Kampen diagram over $F(\mathcal{A})$ (as distinct from a van Kampen diagram over H as above) on Σ_U with boundary label U, thicken the 2-cell D to obtain a disk with an interior region D_0 labelled U, as in Figure 2.4. Remove the interior of D_0 from Σ_U leaving a van Kampen diagram, Δ_U, over $F(\mathcal{A})$ on a surface of genus g, with one boundary component labelled U. It follows from [7] that $\text{genus}_{F(\mathcal{A})}(U) = g$.

Example 2.6. Suppose that we have the quadratic orientable word $U = ABCA^{-1}B^{-1}C^{-1}$ we construct Γ_U and Δ_U as shown in Figure 2.4.



Figure 2.4: Constructing Γ_U and Δ_U

This construction may be carried out in the same way starting with a quadratic t-tuple of words $U_1, \ldots, U_t$. This time let U be the quadratic word $U = U_1 \cdots , U_t$, label the boundary of D with U, as before, define $\Sigma_{U_1,\ldots,U_t}$ to be Σ_U and the associated graph $\Gamma_{U_1,\ldots,U_t}$ to be Γ_U. To construct a van Kampen diagram on a surface with t boundary components labelled U_1, ..., U_t take t letters, X_1, ..., X_t, of $\mathcal{A}\backslash(\cup_{i=1}^{t} \operatorname{supp}(U_i))$, and let $V = X_1^{-1}U_1X_1 \cdots X_t^{-1}U_tX_t$. As before let D_0 be a disk embedded in $\Sigma_{U_1,\ldots,U_t}$ and not meeting $\Gamma_{U_1,\ldots,U_t}$. Label the boundary of D_0 with V and join the ends of corresponding letters of U_i on the boundaries of D and D_0 with disjoint, properly embedded arcs, labelled 1. Remove the interior of D_0 and identify edges labelled X_i with each other, respecting orientation. (See Example 2.7.) The result is the required van Kampen diagram $\Delta_{U_1,\ldots,U_t}$. It follows once more, from [24][Sections 2.3 and 2.4], that $\text{genus}_{F(\mathcal{A})}(U_1, \ldots, U_t) = \text{genus}(\Sigma_{U_1,\ldots,U_t})$.

Example 2.7. Suppose that we have the quadratic orientable pair of words U_1, U_2, where $U_1 = AB$, and $U_2 = CA^{-1}B^{-1}C^{-1}$. Then $U = ABCA^{-1}B^{-1}C^{-1}$ as in Example 2.6. We construct $\Sigma_{U_1,U_2} = \Sigma_U$ and $\Gamma_{U_1,U_2} = \Gamma_U$ as in shown in Figure 2.4. The van Kampen diagram Δ_{U_1,U_2} is obtained by identifying the edges around the outside hexagon on the left of Figure 2.5, as well as those labelled X_i on the inside decagon. The result is a van Kampen diagram on the torus, with two boundary components, labelled AB and $CA^{-1}B^{-1}C^{-1}$.



Figure 2.5: $U_1 = AB$, $U_2 = CA^{-1}B^{-1}C^{-1}$ and Δ_{U_1,U_2} is obtained from the right hand diagram by identifying edges on the outside boundary according to labels.

Let Γ be any directed, connected graph with signature σ. An *Eulerian circuit* in Γ is a circuit C such that,

- C traverses every edge of Γ exactly twice; and
- for all edges e, if $\sigma(e) = (\varepsilon, \delta)$ then (some cyclic permutation of) the edge sequence of C contains the subsequence $e^{\varepsilon}, e^{\delta}$.

By construction the graph Γ_U of a quadratic word U contains an Eulerian circuit, namely the word U, and a similar statement applies to a quadratic tuple of words.

Let Γ be any graph with an Eulerian circuit, T say. Write T around the boundary of a disk and identify the edges, respecting orientation, to obtain a surface S. Then the Euler characteristic $\chi(S)$ of S is given by the formula $v - e + 1$, where v and e are the number of vertices and edges, respectively, of Γ and $\chi(S)$ is equal to $2 - 2\,\text{genus}(S)$. We define genus(Γ), the *genus* of Γ, to be equal to genus(S): that is

$$\text{genus}(\Gamma) = \frac{1 - v + e}{2}. \tag{2.6}$$

As noted above, from [24], if $U_1, \dots, U_t$ is a quadratic t-tuple of words over $F(\mathcal{A})$ then $\operatorname{genus}_{F(\mathcal{A})}(U_1, \dots, U_t) = \operatorname{genus}(\Sigma_{U_1,\dots,U_t})$, so in addition $\operatorname{genus}_{F(\mathcal{A})}(U_1, \dots, U_t) = \operatorname{genus}(\Gamma_{U_1,\dots,U_t})$.

Definition 2.8. *A quadratic t-tuple $w_1, \dots, w_t$ of words is said to be* redundant *if there are x and y in $(\cup_{i=1}^t \operatorname{supp}(w_i))^{\pm 1}$ and disjoint (no overlap) subwords s_1 and s_2 of the cyclic words w_i and w_j, $1 \le i \le j \le t$, such that $s_1 = xy$ and $s_2 = (xy)^{\pm 1}$. Otherwise $w_1, \dots, w_t$ is said to be* irredundant.

Definition 2.9 (Wicks Form)**.** *Let W be a quadratic word in $F(\mathcal{A})$. Then W is called a* Wicks form over $\mathcal{A}$ *if the following conditions hold.*

(i) W is freely cyclically reduced and

(ii) W is irredundant.

Let H be a group with presentation $\langle X | S \rangle$. A (monoid) homomorphism λ from $(\mathcal{A} \cup \mathcal{A}^{-1})^*$ to $F(X)$, such that $\lambda(a^{-1}) =_H \lambda(a)^{-1}$, is called a *labelling function.* If H is the free group on X and h is an element of H, of genus g, then, as shown in [7], there is a Wicks form $W = A_1 \dots A_n$ over $\mathcal{A}$, and a labelling function ψ, such that h is conjugate to $\psi(W)$ and $\psi(W) = \psi(A_1) \cdots \psi(A_n)$ is reduced as written. Indeed Culler (*loc. cit.*) uses this fact to give a description of all solutions of either equation (2.4) or (2.5), when $t = 1$ and $G = F(X)$.

If W is a Wicks form of genus not equal to $1/2$ then the graph Γ_W, associated to W, contains no vertices of degree 1 or 2 (as if it did then either condition (i) or (ii), in the definition of a Wicks form, would be violated). There is only one Wicks form of genus $1/2$, namely $W = X^2$, and it has graph Γ_W consisting of a single vertex and a single edge, lying on the projective plane Σ_W. Consequently, there is a bound on the maximal length of a Wicks form of genus n, given explicitly in the following lemma of M. Culler [7].

Lemma 2.10 ([7, Theorem 3.1])**.** *Let V be a Wicks form over $\mathcal{A}$ such that* $\operatorname{genus}_{F(\mathcal{A})}(V) = m > \frac{1}{2}$*. Then the length of V is at most* $12m - 6$*.*

Lemma 2.11. *Let $h_1, \dots, h_t$ be elements of $F(X)$, $t \ge 1$. Then* $\operatorname{genus}_H(h_1, \dots, h_t) \le m$ *if and only if there exists a Wicks form V over $\mathcal{A}$, of genus m, and a labelling function ψ such that some product $r_1 h_1 r_1^{-1} \cdots r_t h_t r_t^{-1}$, with $r_i \in F(X)$, is equal in H to $\psi(V)$.*

Proof. Suppose that $\operatorname{genus}_H(h_1, \dots, h_t) = g$. Then there exist elements $r_j \in H$, $j = 1, \dots, t$, and either (i) elements $a_i, b_i \in H$, $i = 1, \dots, g$, such that $\prod_{j=1}^t r_j h_j r_j^{-1} = \prod_{i=1}^g [a_i, b_i]$; or (ii) elements $c_i \in H$, such that $\prod_{j=1}^t r_j h_j r_j^{-1} = \prod_{i=1}^{2g} c_i^2$. Let $A_i, B_i, C_i \in \mathcal{A}$. In case (i), for all integers $m \ge g$, $V = \prod_{i=1}^m [A_i, B_i]$ is a Wicks form, of genus m. Define ψ by $\psi(A_i) = a_i$, $\psi(B_i) = b_i$, for $i = 1, \dots, g$, and $\psi(X) = 1$, if $X \in \mathcal{A}$, $X \ne A_i$ or B_i, with $1 \le i \le g$. Then $\psi(V) = \prod_{j=1}^t r_j h_j r_j^{-1}$, as required. In case (ii), for all integers $m \ge 2g$, $V = \prod_{i=1}^m C_i^2$ is a Wicks form, of genus $m/2$. Define ψ by $\psi(C_i) = c_i$, for $i = 1, \dots, 2g$, and $\psi(X) = 1$, if $X \in \mathcal{A}$, $X \ne C_i$, with $1 \le i \le 2g$. Again $\psi(V) = \prod_{j=1}^t r_j h_j r_j^{-1}$, as required.

Conversely, suppose that V is a Wicks form of genus m, that ψ is a labelling function and that $\prod_{j=1}^{t} r_j h_j r_j^{-1}$ is equal in H to $\psi(V)$, for some $r_j \in H$. Without loss of generality, we may assume that h_j is freely cyclically reduced. As $\prod_{j=1}^{t} r_j h_j r_j^{-1}$ is equal to $\psi(V)$ there is a van Kampen diagram D_1, over H, on a surface of genus 0, with $t+1$ boundary components $\beta_1, \dots, \beta_t, \beta$, with β_j labelled h_j and β, labelled $\psi(V)$. Moreover, as in Section 2.3, there is a van Kampen diagram Δ_V over $F(\mathcal{A})$, on a surface Σ_V, of genus m, with one boundary component β', labelled V. Relabelling Δ_V, using the labelling function ψ, we may construct a van Kampen diagram over $F(X)$ with boundary label $\psi(V)$. In more detail: every non-trivial region R of Δ_V has boundary label a cyclic word $a \cdot 1 \cdot a^{-1} \cdot 1$, for some $a \in \mathcal{A}$. Divide the edge labelled a into $|\psi(a)|$ segments and label the resulting edge path $\psi(a)$. The region R can now also be divided into $\psi(a)$ parallel regions, each with boundary label $x \cdot 1 \cdot x^{-1} \cdot 1$, for some $x \in \operatorname{supp}(\psi(a))$. Repeating over all edges and regions results in a van Kampen diagram, over $F(X)$, on Σ_V, with boundary component β', labelled $\psi(V)$. As this is a diagram over $F(X)$ it is *a fortiori* a diagram over H. Thus, attaching D_1 to D_2 by gluing the boundary components β and β' together, using their labels, we obtain a van Kampen diagram over H, on a surface of genus m, with boundary label h. Hence $\operatorname{genus}_H(h_1, \dots, h_t) \leq m$. $\square$

2.5 Extension of a quadratic word over a Hyperbolic group H

Let U be a non-empty quadratic word of genus k (not necessarily reduced or irredundant) over the countably infinite alphabet $\mathcal{A}$ and let Γ_U be its associated graph on the surface Σ_U. Then Γ_U has genus k and signature σ, and its edges are labelled with letters of U. It is notationally convenient to identify a directed edge e of Γ_U with its label $\lambda(A) \in \mathcal{A}^{\pm 1}$ (and e^{-1} with $\lambda(A)^{-1}$) and we shall do so from now on. Thus U itself is an Eulerian circuit of Γ_U. Let $H = \langle X | S \rangle$ be a hyperbolic group in which geodesic triangles in $\Gamma_X(H)$ are δ-thin. We now describe a procedure which, when applied to Γ_U, results in a new graph called an *extension* of the quadratic word U over H. This will be carried out in three steps, but first we need to set up notation.

Roughly speaking, we define the star of a vertex v of Γ_U to be a sufficiently small closed disk, on Σ_U, containing v. To make this precise, let K be the second barycentric subdivision of the cell complex consisting of Γ_U and D on Σ_U. The *star* of v is the convex-hull of the star of v as a 0-cell of the simplicial complex K (the subcomplex of K consisting of all simplices meeting v, together with their faces). As Σ_U is a surface the star of v is a disk and we define the *link* of v to be the boundary of this disk. Then we may view $\operatorname{lk}(v)$ as a cycle graph, with vertices the points where it meets edges of Γ_U (necessarily incident to v) and edges the closures of connected components of $\operatorname{lk}(v) \backslash E(\Gamma_U)$. Let x be a point of intersection of $\operatorname{lk}(v)$ and an edge e of Γ_U. If $v = \tau(e)$ (and $[x, v]$ is contained in the star of v) then we denote the vertex x of $\operatorname{lk}(v)$ by e and if $v = \iota(e)$ we denote x by e^{-1}. (When e is a loop then both e and e^{-1} are vertices of $\operatorname{lk}(v)$.) Thus the union, over all vertices v of Γ_U, of the graphs $\operatorname{lk}(v)$ is isomorphic to the star graph of the word U (the graph with vertices $A^{\pm 1}$, for all $A \in \operatorname{supp}(U)$, and an edge joining A^{ε} to $B^{-\delta}$ for each subword $A^{\varepsilon} B^{\delta}$

of the cyclic word U).

Suppose that a vertex v of Γ_U has degree d. Fix an orientation of v and renumber the edges incident to v (temporarily, for the current purposes) so that the link of v, read according to the chosen orientation, has vertex sequence $e_1^{\varepsilon_1}, e_2^{\varepsilon_2}, \dots, e_d^{\varepsilon_d}$. Then, for all i, the cyclic word U contains the subword $e_i^{\varepsilon_i} e_{i+1}^{-\varepsilon_{i+1}}$ or $e_{i+1}^{\varepsilon_{i+1}} e_i^{-\varepsilon_i}$.

Definition 2.12. *Let v be a vertex of Γ_U, with a chosen orientation, and assume that the link of v has vertex sequence $e_1^{\varepsilon_1}, e_2^{\varepsilon_2}, \dots, e_d^{\varepsilon_d}$. Define the* incidence sequence of v *to be $O_1(v), \dots, O_d(v)$, where*

$$O_1(v) = \begin{cases} 1 & \text{if } U \text{ contains } e_1^{\varepsilon_1} e_2^{-\varepsilon_2} \\ -1 & \text{otherwise} \end{cases}$$

and

$$O_{i+1}(v) = O_i(v) o(e_{i+1}),$$

for $i = 2, \dots d$ (see Definition 2.4).

When the vertex v is fixed we write O_q instead of $O_q(v)$. Thus U contains $(e_i^{\varepsilon_i} e_{i+1}^{-\varepsilon_{i+1}})^{O_i}$, for $i = 1, \dots, d$. It follows that v is incident to an even number of non-alternating edges. Hence, once the orientation of v has been fixed and the first edge e_1 chosen, the incidence sequence is both well-defined and uniquely determined. (Note that, in the case U is orientable, $o(e_i) = 1$, for all i, so $O_1 = \dots = O_d$ and the orientation of v may be chosen so that $O_i = 1$, for all i.)

To keep track of the orientation of letters we define book-keeping functions: μ and ν, from $\{\pm 1\}$ to $\{1, 2\}$, and l and r, from $\{\pm 1\}^2$ to $\{\pm 1\}$, as follows.

$$\begin{array}{ll} \mu(1) = 1, & \mu(-1) = 2 \\ \nu(1) = 2, & \nu(-1) = 1 \\ l(x, y) = \nu(xy), & r(x, y) = \mu(xy). \end{array}$$

Definition 2.13. *Assume v has link with vertex sequence $e_1^{\varepsilon_1}, e_2^{\varepsilon_2}, \dots, e_d^{\varepsilon_d}$, as in the previous definition. Define $\mu_q = \mu_q(v)$, $\nu_q = \nu_q(v)$, $l_q = l_q(v)$ and $r_q = r_q(v)$ as follows, for $q = 1, \dots, d$.*

(i) If $o(e_q) = 1$ then

$$\mu_q = \mu(\varepsilon_q), \nu_q = \nu(\varepsilon_q), l_q = l(O_q, \varepsilon_q) \text{ and } r_q = r(O_q, \varepsilon_q).$$

(ii) Assume that $o(e_q) = -1$ and $\sigma(e_q) = (\varepsilon, \varepsilon)$. In this case $O_{q-1} = -O_q$.

  (a) If the first occurrence in U of the letter e_q^{ε} is the one occurring in the subword $(e_{q-1}^{\varepsilon_{q-1}} e_q^{-\varepsilon_q})^{-O_q}$ then $l_q = 1$ and $r_q = 2$.

  (b) Otherwise the first occurrence in U of the letter e_q^{ε} is the one occurring in the subword $(e_q^{\varepsilon_q} e_{q+1}^{-\varepsilon_{q+1}})^{O_q}$ and $l_q = 2$, $r_q = 1$.

(c) If $O_q = 1$ *then* $\mu_q = r_q$ *and* $\nu_q = l_q$.

(d) Otherwise $\mu_q = l_q$ *and* $\nu_q = r_q$.

(If $d = 1$ *then* e_1 *is necessarily alternating;* $o(e_1) = -1$*, and* O_1 *is* 1 *if and only if the chosen orientation is that induced by reading* U*. In this case* $l_1 = \mu(O_1)$, $r_1 = \nu(O_1)$*, with* μ_1 *and* ν_1 *as in (ii)c and (ii)d above).*

Example 2.14. Let $U = ABA^{-1}B$, so Γ_U is a graph with one vertex and two edges, labelled A and B, on the Klein bottle. We have $\sigma(A) = (1, -1)$, $\sigma(B) = (1, 1)$, $o(A) = 1$ and $o(B) = -1$. The star of the vertex v intersects Γ_U as shown in Figure 2.6(a). Orienting anti-clockwise, v has vertex sequence A^{-1}, B^{-1}, A, B. The following table shows the values of the functions of Definition 2.13 in this case.



(a) (b) (c)

Figure 2.6

q	1	2	3	4
e_q	A	B	A	B
ε_q	-1	-1	1	1
$o(e_q)$	1	-1	1	-1
O_q	1	-1	-1	1
u_q	2	2	1	2
v_q	1	1	2	1
l_q	1	2	1	1
r_q	2	1	2	2

For non-orientable words, in order to construct an extension of U we first fix a particular representative U of $[U]$. For each vertex of Γ_U we also fix an orientation and initial incident edge, so that the incidence sequence is defined, for all vertices.

Step 1. Let e be a directed edge in Γ_U with $u = \iota(e)$ and $v = \tau(e)$ (note that u may be the same vertex as v). In this step we replace e by two new edges e_1 and e_2, where e_1, e_2 are in $(\mathcal{A}\setminus \operatorname{supp}(U))$, such that $\iota(e_1) = \iota(e_2) = u$ and $\tau(e_1) = \tau(e_2) = v$; as follows. As in Section 2.3, consider a disk Δ with boundary divided into $|U|$ consecutive, directed intervals, $I_1, \ldots, I_{|U|}$, labelled with elements of $\mathrm{supp}(U)$, such that, read in a clockwise direction, from the point $I_1 \cap I_{|U|}$, the boundary label of Δ is U. Let $a_1, \ldots, a_{|U|}$ be properly embedded directed arcs in Δ, such that $\iota(a_i) = \iota(I_i)$ and $\tau(a_i) = \tau(I_i)$; and the interiors of a_i and a_j are disjoint, for $i \neq j$. There are precisely two indices i, j such that I_i and I_j, read with their given directions, are labelled e. Assume first that $o(e) = 1$. In this case one of I_i, I_j is oriented clockwise and the other anti-clockwise. If I_i is directed clockwise then the directed arc a_i is labelled e_1 and a_j is labelled e_2. Otherwise a_i is labelled e_2 and a_j labelled e_1. (See Figure 2.7(a).) Now suppose that $o(e) = -1$. If $i < j$ then a_i is labelled with e_1. Otherwise a_i is labelled e_2. (See Figure 2.7(b).) We do this for every edge in Γ_U and form a new graph Γ'_U on Σ_U by deleting the edges of Γ_U. We denote by U' the word obtained by reading the label of a_i instead of that of I_i, when reading round the boundary of Δ, and note that U' is a circuit in Γ'_U. For this circuit we have the following lemma.



(a) (b)

Figure 2.7

Lemma 2.15. *Let v be a vertex of Γ_U with link which has vertex sequence $e_1^{\varepsilon_1}, \ldots, e_d^{\varepsilon_d}$ and incidence sequence $O_1, \ldots, O_d$. Then v is a vertex of Γ'_U of degree $2d$, the link of v in Γ'_U has vertex sequence $e_{1,l_1}^{\varepsilon_1}, e_{1,r_1}^{\varepsilon_1}, \ldots, e_{d,l_d}^{\varepsilon_d}, e_{d,r_d}^{\varepsilon_d}$ and (the cyclic word) U' contains the subwords $(e_{q,r_q}^{\varepsilon_q} e_{q+1,l_{q+1}}^{-\varepsilon_{q+1}})^{O_q}$, for $1 \leq q \leq d$. (See Figure 2.8(a) and (b).)*

Proof. This follows from the definition and the fact that U contains $(e_q^{\varepsilon_q} e_{q+1}^{-\varepsilon_{q+1}})^{O_q}$. ☐

Example 2.16. Applying Step 1 to the word U of example 2.14 we construct the graph shown in Figure 2.6(b); where edges labelled A and B are identified. Deletion of edges A and B then results in a graph $\Gamma_{U'}$ with one vertex v, the link of which intersects $\Gamma_{U'}$ as shown in Figure 2.6(c). Here $U' = A_1 B_1 A_2^{-1} B_2$.

Step 2. In this step we replace vertices of Γ'_U by cycle graphs, as follows. For each vertex v of Γ_U choose a subset $\mathcal{A}_v$ of $\mathcal{A}$ such that

- $|\mathcal{A}_v| = \deg_{\Gamma_U}(v)$,

- $\mathcal{A}_u \cap \mathcal{A}_v = \emptyset$, if $u \neq v$, and
- $\mathcal{A}_v \cap (\mathrm{supp}(U) \cup \mathrm{supp}(U')) = \emptyset$.

For each v, choose a cyclic word $C_v = c_1 \cdots c_d$, where $d = \deg_{\Gamma_U}(v)$ and $\mathcal{A}_v = \{c_1, \ldots, c_d\}$. We shall regard C_v either as a cyclic word or as a directed, labelled, cycle graph, with edges c_i, as expedient. Now let v be a vertex of Γ_U of degree d, with link which has vertex sequence $e_1^{\varepsilon_1}, \ldots, e_d^{\varepsilon_d}$ and incidence sequence $O_1, \ldots, O_d$. Then the link of v in Γ'_U has vertex sequence $e_{1,l_1}^{\varepsilon_1}, e_{1,r_1}^{\varepsilon_1}, \ldots, e_{d,l_d}^{\varepsilon_d}, e_{d,r_d}^{\varepsilon_d}$. Now assume that the cycle C_v is $c_1 \cdots c_d$, with vertices $\iota(c_i) = v_i$ and $\tau(c_i) = v_{i+1}$, subscripts modulo d. Remove the vertex v from Γ'_U and replace it with C_v; according to the following scheme. If $d \geq 2$ then

$$\begin{array}{rrcll} & \tau(e_{d,r_d}^{\varepsilon_d}) = \tau(e_{1,l_1}^{\varepsilon_1}) & = & v_1 & \\ \text{and} & \tau(e_{q-1,r_{q-1}}^{\varepsilon_{q-1}}) = \tau(e_{q,l_q}^{\varepsilon_q}) & = & v_q, & \text{for } q = 2, \ldots, d. \end{array}$$

See Figure 2.8. If v is a vertex of Γ_U of degree one, then the edge incident to v must be alternating and U must contain the subword $e^{\varepsilon} e^{-\varepsilon}$, $\varepsilon = \pm 1$. In this case add a loop with the label c_1 to v, as in Figure 2.9. On the surface Σ_U we require that this loop lies in the disk which is bounded by e_1 and e_2 and which meets no other edge of Γ'_U, as illustrated. In all cases we say that v has been *extended by* C_v. After each vertex has been extended we obtain a new graph which we shall call Γ''_U. We can still read the closed path U' in this graph. In fact it is now a Hamiltonian cycle. We shall call U' the *Hamiltonian cycle associated with* U.

Step 3. In this step we choose a labelling function for the new graph, which allows us (eventually) to regard it as a van Kampen diagram over H. Consider a directed edge e of the original graph Γ_U. In Step 1 this was replaced by a pair of edges (e_1, e_2). Then in Step 2 we extended the end points of these edges to a subgraph of Γ''_U of the form shown in Figure 2.10, where x and y are letters occurring in $C_{\iota(e)}$ and $C_{\tau(e)}$, respectively. Let $\psi : (\mathcal{A} \cup \mathcal{A}^{-1})^* \to F(X)$ be a labelling function such that

(a) $\psi(a)$ is H-minimal, for all $a \in \mathcal{A}$,

(b) $\psi(e_1) =_H \psi(x)\psi(e_2)\psi(y)$, for all pairs of directed edges (e_1, e_2) corresponding to an edge e of Γ_U, with x and y as in Figure 2.10, and

(c) if $U' = a_1 \cdots a_r$ is the Hamiltonian cycle associated to U, where $a_i \in \mathcal{A}^{\pm 1}$, then $\psi(a_1) \cdots \psi(a_r)$ is freely cyclically reduced (as written).

Then the resulting pair (Γ''_U, ψ) is called an *extension of* U *over the group* H.

2.6 Genus and length of an extension of U over H

Suppose that we have an extension (Γ''_U, ψ) of a non-empty quadratic word U of genus k. We define the *length of the extension* to be the sum of the lengths, in $F(X)$, of the labels of the cyclic words added in Step 2 of the extension. That is, if $v_1, v_2, \ldots, v_m$ are the vertices


$e_1^{\varepsilon_1}$
$e_d^{\varepsilon_d}$
$e_2^{\varepsilon_2}$
$e_3^{\varepsilon_3}$
v
$e_{d-1}^{\varepsilon_{d-1}}$
$\longrightarrow$
$e_{1,l_1}^{\varepsilon_1}$
$e_{1,r_1}^{\varepsilon_1}$
$e_{d,r_d}^{\varepsilon_d}$
$e_{d,l_d}^{\varepsilon_d}$
$e_{2,l_2}^{\varepsilon_2}$
$e_{2,r_2}^{\varepsilon_2}$
$e_{d-1,r_{d-1}}^{\varepsilon_{d-1}}$
$e_{d-1,l_{d-1}}^{\varepsilon_{d-1}}$
$e_{3,l_3}^{\varepsilon_3}$
$e_{3,r_3}^{\varepsilon_3}$
v
$\swarrow$
c_1
c_2
c_d
v_1
v_2
v_3
v_d

(a)

(b)

(c)

Figure 2.8: $o(e_1) = 1$, $o(e_2) = o(e_d) = -1$, $O_1 = 1$


$e_{\mu(\varepsilon)}^{\varepsilon}$
$e_{\nu(\varepsilon)}^{\varepsilon}$
$e_{\mu(\varepsilon)}^{\varepsilon}$
c_1
$e_{\nu(\varepsilon)}^{\varepsilon}$

Figure 2.9



Figure 2.10

of Γ_U, and these are extended by cyclic words $C_1, C_2, \ldots, C_m$ respectively, then the length of the extension is

$$\sum_{i=1}^{m} |\psi(C_i)|.$$

Definition 2.17. *Let* (Γ''_U, ψ) *be an extension of* U *over* H*, let* $v_1, \ldots, v_t$ *be a subset of the vertices of* Γ_U *such that* v_i *is extended by a cyclic word* C_i *and let* $w_i = \psi(C_i)$*, for* $i = 1, \ldots, t$*. Then we say that* (Γ''_U, ψ) *is a* genus g joint extension on $(v_1, \ldots, v_t)$ by words $(w_1, \ldots, w_t)$*, if*

- *(i)* $g = \text{genus}_H(w_1, \ldots, w_t) + t - 1$ *and*
- *(ii) if* $t = 1$ *and* v_1 *has degree* 1 *or* 2 *then either*
  - *(a)* $g \geq \frac{1}{2}$ *or*
  - *(b)* $U = A^{\pm 2}$*, for some* $A \in \mathcal{A}$*.*

If $\text{genus}^+_H(w_1, \ldots, w_t) = g - t + 1$*, in (i) above, then we say that* (Γ''_U, ψ) *is an* orientable *extension on* $(v_1, \ldots, v_t)$ *and otherwise that it is* non-orientable *on* $(v_1, \ldots, v_t)$*.*

(Note that condition (ii) implies that if $U \neq A^{\pm 2}$ and $g = 0$, which means that t must be 1, then v_1 has degree at least 3.)

Definition 2.18. *Partition the vertices of* Γ_U *into* p *sets* $V_1, \ldots, V_p$*. If*

- *(i)* (Γ''_U, ψ) *is a genus* g_i *joint extension on the vertices in the set* V_i*, for all* $i = 1, \ldots, p$*, and*
- *(ii)* $\sum_{i=1}^{p} g_i = g$*,*

then (Γ''_U, ψ) *is called a* genus g extension of U over H*. If* U *is orientable and* (Γ''_U, ψ) *is orientable on* V_i*, for* $1 \leq i \leq p$*, then this extension is called* orientable*. Otherwise it is said to be* non-orientable*.*

Example 2.19. (A genus 3 extension)
Consider the quadratic orientable word $U = ABCC^{-1}B^{-1}A^{-1}$ of genus 0. Let u_1, u_2, v_1, v_2 be the vertices of the associated graph Γ_U, as shown in Figure 2.11. We construct a genus 2 joint extension on the vertices u_1 and u_2, by words w_1 and w_2 respectively, and a genus 1 joint extension on the vertices $v_1$and v_2, by words z_1 and z_2 respectively (see Figure 2.11). Here all words in the generators of H which are mentioned are assumed to be H-minimal, $w_2 = w_{21}w_{22}$ and $z_1 = z_{11}z_{12}$. The labelling function ψ is such that $\psi(c_1) = w_1$, $\psi(d_2) = z_2$, $\psi(c_{ij}) = w_{ij}$, $\psi(d_{ij}) = z_{ij}$,



Figure 2.11

$$\begin{aligned}
\psi(A_1) &=_H w_1\psi(A_2)w_{22},\\
\psi(B_1) &=_H w_{21}\psi(B_2)z_{12},\\
\psi(C_1) &=_H z_{11}\psi(C_2)z_2,
\end{aligned}$$

$\psi(A_1)\psi(B_1)\psi(C_1)\psi(C_2^{-1})\psi(B_2^{-1})\psi(A_2^{-1})$ is freely cyclically reduced, w_1w_2 is a commutator in H and $z_1z_2 = 1$ in H.

Lemma 2.20. *Let (Γ''_U, ψ) be a genus g extension of the quadratic word U of genus k and let U' be the Hamiltonian cycle associated to U. Let $h_1, \dots, h_s$ be elements of H such that $\prod_{j=1}^{t} r_jh_jr_j^{-1} =_H \psi(U')$, for some $r_j \in H$. If (Γ''_U, ψ) is orientable, then* $\text{genus}^+_H(h_1, \dots, h_s) \le g + k$ *and otherwise* $\text{genus}^-_H(h_1, \dots, h_s) \le g + k$.

Proof. The graphs Γ_U and Γ''_U are embedded on a closed surface Σ_U, by construction. As Σ_U is constructed by identifying segments of the boundary of the disk D, the Hamiltonian circuit U', associated to U, labels a closed path in Γ''_U which bounds an open disk embedded in Σ_U and not meeting Γ_U. Remove this disk from Σ_U, to leave a closed surface $\Sigma_{U,0}$, with one boundary competent β_0. Then $\Sigma_{U,0}$ is the convex-hull of the 2-complex, which has 1-skeleton $\Gamma_{U''}$ and 2-cells the closures of components of $\Sigma_{U,0}\backslash\Gamma_{U''}$, and moreover β_0 is labelled $\psi(U')$. As $\prod_{j=1}^{s} r_jh_jr_j^{-1} =_H \psi(U')$ there exists a van Kampen diagram Δ_H, over H, on surface of genus 0, with $s + 1$ boundary components $\beta_1, \dots, \beta_s, \beta$, with β_j labelled h_j and β labelled $\psi(U')$. Identifying β' with β_0, along $\psi(U')$, results in a surface Σ'_U, with

s boundary components $\beta_1, \dots, \beta_s$, labelled with the h_j's. Indeed we may regard Σ'_U as obtained from Σ_U by removing the interior of s disks. Moreover Γ''_U is embedded in Σ'_U and does not meet β.

Now assume the vertices of Γ_U were $v_1, \dots, v_t$ and assume that v_i has been extended by the cycle graph C_i. For each i, the cycle C_i bounds a disk D_i in Σ'_U, the interior of which does not meet Γ''_U. For all i, remove the interior of D_i from Σ'_U to leave a surface Σ''_U with $t+s$ boundary components labelled $h_1, \dots, h_s$ and $\psi(C_1), \dots, \psi(C_t)$. From the conditions imposed on the labelling function ψ it follows that the labelled graph Γ''_U, together with the diagram Δ_H, form a van Kampen diagram over H, on the surface Σ''_U.

Now suppose that the vertices of Γ_U are partitioned into sets V_1, ..., V_p in such a way that (Γ''_U, ψ) forms a genus g_i joint extension on V_i, with $\sum_{i=1}^{p} g_i = g$. Let $w_i = \psi(C_i)$, for all i. Then, if $V_i = \{v_{i,1} \dots, v_{i,t_i}\}$, by definition we have $\text{genus}_H(w_{i,1} \dots, w_{i,t_i}) = g_i - t_i + 1$. Hence, there exists a van Kampen diagram, over H, on a compact surface T_i of genus $g_i - t_i + 1$, with t_i boundary components labelled $w_{i,1} \dots, w_{i,t_i}$. Attach the boundary component of T_i labelled $w_{i,j}$ to the boundary component of Σ''_U labelled $w_{i,j}$, for $j = 1, \dots, t_i$. In this way we attach a handle, or cross-cap, of genus $t_i - 1 + \text{genus}(T_i) = g_i$ to Σ''_U. Repeat the process for all sets V_i of the partition of the vertices of Γ_U. As Σ_U has genus k, so does Σ''_U, so the result is a surface Σ of genus $k + \sum_{i=1}^{p} g_i = k + g$. We have now constructed a van Kampen diagram, over H, on Σ, with s boundary components, labelled $h_1, \dots, h_s$. Hence $\text{genus}_H(h_1, \dots, h_s) \le g + k$. ☐

Example 2.21. In Example 2.19 we constructed a genus 3 extension consisting of a joint genus 2 extension by words w_1, w_2, and a joint genus 1 extension by z_1 and z_2. To construct the corresponding van Kampen diagram on a surface of genus 3 first embed the graph Γ'' in the disk, with boundary labelled $A_1B_1C_1C_2^{-1}B_2^{-1}A_2^{-1}$. Then remove the interior of the disks labelled c_1, $c_{21}c_{22}$, $d_{11}d_{12}$ and d_2. Finally, attach a torus with two boundary components, labelled c_1 and $c_{21}c_{22}$ and a sphere with two boundary components labelled d_2 and $d_{11}d_{12}$, in the obvious way: see Figure 2.12.

2.7 The main theorems

We are now in a position to state the main results of this paper, which give a method of constructing forms for elements of (orientable or non-orientable) genus n in H.

Definition 2.22. *Let H be a δ-hyperbolic group, with respect to the generating set X. For an integer $k \ge 0$, let $B_H(k)$ be the set of elements of H represented by words of length at most k in $F(X)$. Define $M = |B_H(4\delta)|$: and, for a positive element $n \in \mathbb{Z}[\frac{1}{2}]$, define $K(n) = 12n - 6$, $n \ge 1$, $K(1/2) = 2$, $K(0) = 0$ and $l(n) = \delta(\log_2(K(n)) + 1)$.*

Assume that $h \in H$, that the orientable genus of h is $\text{genus}^+_H(h) = g$, that U is an orientable Wicks form, over $\mathcal{A}$, of genus g, and that ϕ is a labelling function such that $\phi(U)$ is conjugate to h. Then $\phi(U)$ has orientable genus g in $F(X)$. (It cannot have orientable genus less than g, otherwise h would, as well.) It follows, from [7], that there exists an orientable Wicks form W of genus g over $\mathcal{A}$, and a labelling function ψ such



Figure 2.12: A van Kampen diagram constructed from an extension of U

that $\psi(W)$ is conjugate, in $F(X)$, to $\phi(U)$, and the cyclic word obtained by substituting $\psi(X)$ for X, for all $X \in \mathrm{supp}(W)$, is reduced as written, as a word in $F(X)$. That is $\psi(W)$ is freely cyclically reduced and $|\psi(W)| = 2\sum_{X\in\mathrm{supp}(W)} |\psi(X)|$. Call such a labelling function *cancellation free* on W. The same holds for the non-orientable genus: replacing "orientable" by "non-orientable" in the paragraph above.

Let W be a Wicks form and let D be a proper subset of $\mathrm{supp}(W)$. The word U obtained from W by setting all elements of D equal to 1 is called a *specialisation* of W. For example, up to permutation of letters in the support, the Wicks form $ABCA^{-1}B^{-1}C^{-1}$ has specialisations AA^{-1}, $ABA^{-1}B^{-1}$ and $ABCA^{-1}B^{-1}C^{-1}$.

Theorem 2.23. *Let h be a word in $X \cup X^{-1}$, let n be a positive integer and let $l = l(n)$. Then $\mathrm{genus}^+_H(h) = n$ if and only if n is the minimal integer such that the following holds. There exist words $F, R \in F(X)$ such that $h =_H RFR^{-1}$, and an orientable Wicks form W over $\mathcal{A}$, of genus n, satisfying either 1 or 2 below.*

1. *$F = \theta(W)$, where θ is a cancellation free labelling on W, $|\theta(E)| \le 12l+M+4$, for all $E \in \mathrm{supp}(W)$ (so $|F| \le (12n-6)(12l+M+4)$) and $|R| \le |h|/2+6l+3M/2+2\delta+7/2$.*
2. *F is H-minimal and $F =_H \psi(U')$, where U' is the Hamiltonian cycle of an orientable, genus g extension (Γ''_U, ψ), of length at most $2(12n-6)(12l+M+4)$, of some specialisation U of W, of genus k over $\mathcal{A}$, where $n = g + k$. In this case $|R| \le |h|/2 + 2\delta$.*

Theorem 2.24. *Let h be a word in $X \cup X^{-1}$, let $n \in \mathbb{Z}[\frac{1}{2}]$, $n > 0$ and let $l = l(n)$. Then $\mathrm{genus}^-_H(h) = n$ if and only if n is the minimal positive element of $\mathbb{Z}[\frac{1}{2}]$ such that the following hold. There exist words $F, R \in F(X)$ such that $h =_H RFR^{-1}$, and a non-orientable Wicks form W over $\mathcal{A}$, of genus n, satisfying either 1 or 2 below.*

1. *$F = \theta(W)$, where θ is a cancellation free labelling on W, $|\theta(E)| \le 12l + M + 4$, for all $E \in \mathrm{supp}(W)$; so $|F| \le K(n)(12l + M + 4)$. In this case $|R| \le |h|/2 + 6l + 3M/2 + 2\delta + 7/2$.*
2. *F is H-minimal and $F =_H \psi(U')$, where U' is the Hamiltonian cycle of a non-orientable, genus g extension (Γ''_U, ψ), of length at most $2K(n)(12l + M + 4)$, of some specialisation U of W of genus k over $\mathcal{A}$, where $n = g + k$. In this case $|R| \le |h|/2 + 2\delta$.*

The proof of these two theorems is the content of Section 4.

3 Applications

3.1 Forms for Commutators

We use Theorem 2.23 to obtain a full list of all possible forms for commutators in $H = \langle X|S\rangle$. Since in this case $n = 1$, it follows that $l = \delta(\log_2(6) + 1)$.

Proposition 3.1. *An element $h \in H$ is a commutator if and only if there are words R and F in $F(X)$, such that $h =_H RFR^{-1}$, where F takes one of the following forms, $|R| \leq |h|/2 + 6l + 3M/2 + 2\delta + 7/2$ in case 1, and $|R| \leq |h|/2 + 2\delta$ in cases 2, 3 and 4.*

1. *$F = XYZX^{-1}Y^{-1}Z^{-1}$, for $X, Y, Z \in F(X)$ with $|X|$, $|Y|$, $|Z| \leq 12l + M + 4$.*
2. *$F = A_1A_2^{-1}$ with $A_1 =_H \xi_1^{-1}A_2\xi_2$, where $|\xi_1| + |\xi_2| \leq 12(12l + M + 4)$ and ξ_1 is conjugate to ξ_2 in H.*
3. *$F = A_1B_1A_2^{-1}B_2^{-1}$ with $A_1 =_H \xi_1A_2\xi_3$, $B_1 =_H \xi_4B_2\xi_2$, where $|\xi_1| + |\xi_2| + |\xi_3| + |\xi_4| \leq 12(12l + M + 4)$ and $\xi_1\xi_2\xi_3\xi_4 =_H 1$.*
4. *$F = A_1B_1C_1A_2^{-1}B_2^{-1}C_2^{-1}$ with $A_1 =_H \xi_1A_2\rho_1$, $B_1 =_H \rho_2B_2\xi_2$ and $C_1 =_H \xi_3C_2\rho_3$, where $|\xi_1| + |\xi_2| + |\xi_3| + |\rho_1| + |\rho_2| + |\rho_3| \leq 12(12l + M + 4)$ and $\xi_1\xi_2\xi_3 =_H \rho_1\rho_2\rho_3 =_H 1$.*

Proof. By Theorem 2.23, h is conjugate to a word F which either has form 1 above or is H-minimal and obtained by an orientable, genus g extension, of length at most $12(12l + M + 4)$, of some specialisation U of the Wicks form $ABCA^{-1}B^{-1}C^{-1}$, of genus k, where $g + k = 1$. The bounds on $|R|$ also follow directly from the theorem. There are only three specialisations which can have a suitable extension.

(i) If $k = 0$ and $g = 1$, take a quadratic orientable word $U = AA^{-1}$, of length 2 and genus 0, with a joint genus 1 extension constructed on the two vertices of Γ_U.

(ii) If $k = 1$ and $g = 0$ there are two possibilities.

  (a) The orientable word $U = ABA^{-1}B^{-1}$ of genus 1 with a genus 0 extension constructed on the only vertex of Γ_U.

  (b) The orientable word $U = ABCA^{-1}B^{-1}C^{-1}$ of genus 1 with a genus 0 extension constructed on both of the vertices of Γ_U.

(i) We extend the graph Γ_U as shown in Figure 3.1. By Theorem 2.23, F takes the form of the Hamiltonian cycle $A_1A_2^{-1}$ in the extended graph and from the nature of the extension constructed on U, it is clear that we have form 2.



Figure 3.1

(ii)(a) We extend the graph Γ_U as shown in Figure 3.2. By Theorem 2.23, F takes the form of the Hamiltonian cycle $A_1B_1A_2^{-1}B_2^{-1}$ in the extended graph and from the nature of the extension constructed on U, it is clear that we have form 3.



Figure 3.2



Figure 3.3

(ii)(b) Finally, we extend the graph Γ_U as shown in Figure 3.3. Again, by Theorem 2.23, F takes the form of the Hamiltonian cycle $A_1B_1C_1A_2^{-1}B_2^{-1}C_2^{-1}$ in the extended graph and from the nature of the extension constructed on U, it is clear that we have form 4. Hence h is conjugate to some F which takes one of the required forms.

□

3.2 Forms for Squares

For a list of squares we use Theorem 2.24. In this case $n = 1/2$, so $l = 3\delta$.

Proposition 3.2. *An element $h \in H$ is a square if and only if there are words R and F in $F(X)$, such that $h =_H RFR^{-1}$, where either*

1. $F = X^2$, *for $X \in F(X)$ with $|X| \le 12l+M+4$ and $|R| \le |h|/2+12l+3M/2+2\delta+7/2$; or*
2. $F = A_1A_2$ *with $A_1 =_H \xi A_2 \xi$, where $|\xi| \le 5l + M + 4$ and $|R| \le |h|/2 + 2\delta$.*

Proof. By Theorem 2.24, h is conjugate to a word F which either has form 1 above or is obtained by a non-orientable, genus g extension, of length at most $4(12l + M + 4)$, of some specialisation of the Wicks form A^2, of genus k, where $g + k = 1/2$. There is one possible specialisation which can have a suitable extension, namely the word $U = A^2$, of genus $k = 1/2$. Therefore we have a genus 0 extension over the unique vertex v of Γ_U. We extend the graph Γ_U as shown in Figure 3.4. Following through the proof of Proposition 4.15, we see that the length of this extension is at most $2(5l + M + 4)$, since in this case v is an internal vertex and the words $b_1(A) = a_2(A) = 1$, while $a_1(A) = b_2(A)$ has length at most $5l + M + 4$. Also, as the extension is of genus 0 we have $\xi = \xi_2 =_H \xi_1^{-1}$, and $\xi A_2 \xi A_1^{-1} =_H 1$, as required. □



Figure 3.4

3.3 Solutions of quadratic equations in hyperbolic groups

In this section we extend Theorems 2.23 and 2.24 to apply to quadratic tuples of elements of H. That is we show that, if $\text{genus}_H(c_1, \ldots, c_t) \le g$ then there exist conjugators r_i, linearly bounded in terms of the sum of lengths of the c_i, such that $\prod_{i=1}^{t} c_i^{r_i}$ has one of the forms given in Theorem 2.23 or 2.24, as appropriate. In the process we give bounds on the lengths of images of variables in (minimal) solutions of quadratic equations over hyperbolic groups. These are similar to those found over free groups by Kharlampovich and Vdovina [20] and by Lysenok and Myasnikov [21]. For a torsion-free hyperbolic group H, Kharlampovich, Mohajeri, Taam and Vdovina [19] founds such bounds which are in somewhat different from those we give below. In fact they compute a number N, dependent only H, such that if a quadratic equation Q is solvable in H then there is a solution in which the variables are replaced by words of length at most NL^d, where L is the length of the the equation Q, as a word in the generators of H and the variables of Q, and $d = 3$, if Q is orientable, and 4 otherwise. In contrast the bounds given belown depend on the genus of the equation, but are linear (and torsion is not excluded). As usual, suppose that H is δ-hyperbolic with respect to the presentation $\langle X|S\rangle$. First we use a result from [24] to bound the length of solutions of quadratic equations over H.

Theorem 3.3 (*cf.* [19][Theorem 1], [21][Theorem 1.1])**.** *Let H be a δ-hyperbolic group, with respect to the presentation $\langle X|S\rangle$, let $g \in \mathbb{Z}[\frac{1}{2}]$ and let $c_1, \dots, c_t$ be elements of $F(X)$. Then there exist constants A and B, dependent only on δ, $|X|$, g and t such that the following hold.*

1. *If* $\mathrm{genus}^+(c_1, \dots, c_t)_H \le g$ *then there exists a solution ϕ to equation* (2.4) *such that*

$$|\phi(v_l)|, |\phi(x_i)|, |\phi(y_i)| \le A\sum_{j=1}^{t} |c_j| + B,$$

$i = 1, \dots, g$, $l = 1, \dots, t$.

2. *If* $\mathrm{genus}^-(c_1, \dots, c_t)_H \le g$ *then there exists a solution ϕ to equation* (2.5) *such that*

$$|\phi(v_l)|, |\phi(z_i)| \le A\sum_{j=1}^{t} |c_j| + B,$$

$i = 1, \dots, 2g$, $l = 1, \dots, t$.

Proof. First note that we may assume that R contains elements of length at most 8δ (see for example [1]). Let $c_1, \dots, c_t$ be orientable and let $X_i = \phi(x_i)$, $Y_i = \phi(y_i)$ and $V_l = \phi(v_l)$ be a solution to the equation (2.4). As in Section 2.3 there is a van Kampen diagram $\mathcal{K}$ on a surface Σ of genus g, with boundary components labelled c_i, obtained from a disk with boundary labelled $\prod_{i=1}^{t} V_i^{-1} c_i V_i \prod_{i=1}^{g} [X_i, Y_i]$. We assume that the number of regions of $\mathcal{K}$ is minimal, among all such van Kampen diagrams. From [24][Lemma 11] we may cut $\mathcal{K}$ along edges of 2-cells to obtain a van Kampen diagram on the disk; in such a way that the boundary label of the disk has length at most $\sum_{j=1}^{t} |c_j| + C$, for some constant C dependent only on δ, $|X|$, g and t. As H is hyperbolic it has a linear Dehn function $f(n) = Dn + E$, for some constants D and E, dependent only on δ, so the van Kampen diagram on the disk has at most $D(\sum_{j=1}^{t} |c_j| + C) + E = D\sum_{j=1}^{t} |c_j| + E'$ regions. By construction the same is true of $\mathcal{K}$. Each of X_i, Y_i and V_l on $\mathcal{K}$ corresponds to a simple path on $\mathcal{K}$, formed from a union of boundary components of 2-cells. As the 2-cells of $\mathcal{K}$ have boundaries of length at most 8δ, we may assume that each of V_l, X_i and Y_i has length bounded by $4\delta(D\sum_{j=1}^{t}|c_j|+E') = A\sum_{j=1}^{t}|c_j|+B$, with $A = 4D\delta$, $B = 4E'\delta$. This proves 1 and 2 follows similarly. $\square$

A *system* of equations is a set of equations $(w_i = 1)_{i\in I}$, where I is an indexing set, and a *solution* to this system of equations over H is an H-map $\phi : F(X) * F(\mathcal{A}) \to H$ such that $\phi(w_i) =_H 1$, for all $i \in I$.

Corollary 3.4 (*cf.* [20][Theorem 2])**.** *Let* $\mathbf{w} = w_1, \dots, w_t$ *be a quadratic tuple of words over* $\mathcal{A}$ *of genus* $g \in \mathbb{Z}[\frac{1}{2}]$ *and let* $c_1, \dots, c_t$ *be elements of* $F(X)$ *such that the system of equations* $(w_ic_i = 1)_{i=1}^{t}$ *has solution in* H*. Let* $d = \sum_{i=1}^{t}|c_i|$*. Then there exist constants* A *and* B*, dependent only on* δ*,* $|X|$*,* g *and* t*, such that the following hold.*

1. *If* $\mathbf{w}$ *is orientable then there exists a solution* ϕ *to the equation* (2.4) *over* H *such that* $|\phi(a)| \le Ad + B$*, for all* $a \in \{x_i, y_i \,|\, 1 \le i \le g\} \cup \{v_i \,|\, 1 \le i \le t\}$*.*
2. *If* $\mathbf{w}$ *is non-orientable then there exists a solution* ϕ *to the equation* (2.5) *over* H *such that* $|\phi(a)| \le Ad + B$*, for all* $a \in \{z_i \,|\, 1 \le i \le 2g\} \cup \{v_i \,|\, 1 \le i \le t\}$*.*

Proof. Let ψ be a solution to $(w_ic_i = 1)_{i=1}^{t}$. For each i label the boundary of a disk D_i with $w_iv_i^{-1}c_iv_i$, where $v_i \in \mathcal{A}$, $v_i \notin \mathrm{supp}(w_j)$, for all j, and $v_i \ne v_j$, $i \ne j$. Setting $\psi(v_i) = 1$, for all i, the system of equations $(w_iv_i^{-1}c_iv_i = 1)_{i=1}^{t}$ also has solution ψ. Let D be the disjoint union of these disks. The quotient of D obtained by identifying directed edges, according to their label in $\mathcal{A}$, is a surface Σ (not necessarily connected) with t boundary components. Collapsing the arcs labelled $v_i^{-1}c_iv_i$, on the boundary of each D_i, to a point collapses all boundary components of Σ to a point and, as noted above Definition 2.8, as $\mathbf{w}$ has genus g, the surface obtained from Σ, by this collapse, has genus g. Therefore Σ also has genus g. If $\mathbf{w}$ is orientable then so is Σ and, read with an appropriate orientation of Σ, the the boundary labels of Σ are $c_1, \dots, c_t$ (see Section 2.3). If $\mathbf{w}$ is non-orientable, then with an appropriate choice of orientation of each boundary component, the boundary labels of Σ are again $c_1, \dots, c_t$.

Now relabel each directed edge on D_i by applying ψ: that is an edge labelled $a \in \mathcal{A}$ is relabelled $\psi(a)$. This results in a disk with boundary label $\psi(w_i)c_i =_H 1$, for each i. Hence there is a van Kampen diagram over H on each disk D_i. Thus there is also a van Kampen diagram on Σ. Therefore there is a corresponding solution ϕ to the equation (2.4) in the case where $\mathbf{w}$ is orientable and to (2.5) otherwise. That is $\mathrm{genus}_H^{\pm}(c_1, \dots, c_t) \le g$ and the result follows from Theorem 3.3. $\square$

Theorem 3.5. *With the hypotheses of Theorem 3.3, there exist constants* A' *and* B'*, dependent only on* δ*,* $|X|$*,* g *and* t *such that the following hold. If* $\mathrm{genus}^{\pm}(c_1, \dots, c_t)_H = g$ *there exist* $F, r_1, \dots, r_t \in F(X)$ *such that*

$$\prod_{i=1}^{t} r_i^{-1}c_ir_i = F \quad \text{and} \quad |r_i| \le A'\sum_{j=1}^{t}|c_j| + B',$$

for $i = 1, \dots, t$, where

1. *if* $\mathrm{genus}^+(c_1, \dots, c_t)_H = g$ *then F satisfies 1 or 2 of Theorem 2.23 and*
2. *if* $\mathrm{genus}^-(c_1, \dots, c_t)_H = g$ *then F satisfies 1 or 2 of Theorem 2.24.*

Proof. Let $c_1, \dots, c_t$ be orientable and let ϕ be a solution to equation (2.4). From Theorem 3.3, there exist constants A and B such that $|\phi(a)| \le A \sum_{i=1}^t |c_i| + B$, for all variables a in the equation. Then $h = \prod_{i=1}^t V_i^{-1} c_i V_i$ has orientable genus g, where $V_i = \phi(v_i)$, and $|h|$ is bounded by $(2tA + 1) \sum_{i=1}^t |c_i| + 2tB$. From Theorem 2.23 there exist $R, F \in F(X)$ such that $h = RFR^{-1}$ and F and R satisfy 1 or 2 of the theorem. Here R is bounded by $|h|/2 + C'$, for some constant C' dependent only on δ, $|X|$ and g, so we have $F = \prod_{i=1}^t r_i^{-1} c_i r_i$, where $r_i = R^{-1} V_i R$; and $|r_i| \le A' \sum_{i=1}^t |c_i| + B'$, for constants A' and B' dependent only on δ, $|X|$, g and t. This proves 1 and 2 follows similarly. □

4 Proof of Theorems 2.23 and 2.24

4.1 Decomposition over short and long edges

The proofs of Theorems 2.23 and 2.24 involve consideration of a genus n Wicks form W over $\mathcal{A}$ and a homomorphism θ from $F(\mathcal{A})$ to $F(X)$ such that

- $\theta(W)$ is conjugate to h in H and
- amongst all such Wicks forms (of the appropriate orientation) and maps, W and θ are minimal, in a sense made precise below.

In outline: we shall first establish some preliminary lemmas involving Wicks forms and then show that W and θ either satisfy condition 1 of our theorems, or W can be reduced to a quadratic word U, by setting certain letters equal to 1, and from U an extension satisfying condition 2 may be constructed.

We abbreviate notation to allow discussion of both theorems simultaneously. We write genus_H to mean genus^+_H or genus^-_H, and "genus" instead of "orientable genus" or "non-orientable genus", whenever possible. We omit "orientable" and "non-orientable" when talking of Wicks forms; the context should make it clear which is meant.

Proof of Theorems 2.23 and 2.24. From Lemmas 2.11 and 2.20 it follows that if the conditions of either Theorem hold then $\mathrm{genus}_H(h) = n$. It remains to prove the converse. In the orientable case (Theorem 2.23) we note that, from Lemma 2.11, there exists no genus m Wicks form V, $m < n$, with a labelling function ψ such that $\psi(V)$ is conjugate to h in H. By hypothesis, the genus of h is equal to n in $H = \langle X|S \rangle$. Therefore, in the case of Theorem 2.23, $n \in \mathbb{Z}$ and there exist words $a_i, b_i \in F(X)$, for $i = 1, \dots, n$, such that

$$h =_H [a_1, b_1][a_2, b_2] \cdots [a_n, b_n].$$

The quadratic orientable word $U = [A_1, B_1] \cdots [A_n, B_n]$, over $\mathcal{A}$, is a genus n Wicks form. Let $\phi : F(\mathcal{A}) \to F(X)$ be the labelling function defined by $\phi(A_i) = a_i$ and $\phi(B_j) = b_j$, for all $i, j = 1, \ldots, n$, and $\phi(X) = 1$, otherwise. Then $\phi(U) = h$. Moreover, $\phi(U)$ has genus n in $F(X)$. In this (orientable) case let $\mathcal{F} = \mathcal{F}(h)$ be the set of pairs (U, ϕ) where U is an orientable genus n Wicks form and ϕ is a labelling function such that $\phi(U)$ is conjugate to h in H.

In the non-orientable case, Theorem 2.24, $n \in \mathbb{Z}[1/2]$ and there exist words $c_i \in F(X)$, for $i = 1, \ldots, 2n$, such that

$$h =_H c_1^2 \cdots c_{2n}^2.$$

The non-orientable quadratic word $U = C_1^2 \cdots C_{2n}^2$, over $\mathcal{A}$, is a genus n Wicks form. In this case let $\phi : F(\mathcal{A}) \to F(X)$ be the labelling function given by $\phi(C_i) = c_i$, $i = 1, \ldots, 2n$, and $\phi(X) = 1$, otherwise. Then $\phi(U) = h$ and $\phi(U)$ has genus n in $F(X)$. In this (non-orientable) case, let $\mathcal{F} = \mathcal{F}(h)$ be the set of pairs (U, ϕ) where U is a non-orientable genus n Wicks form and ϕ is a labelling function such that $\phi(U)$ is conjugate to h in H.

In both orientable and non-orientable cases, we call a pair $(W, \theta) \in \mathcal{F}$ *minimal* if $|\theta(W)|$ is minimal amongst all pairs in $\mathcal{F}$. Since we have shown that, in both cases, there is at least one pair in $\mathcal{F}$ there is always a minimal pair. As above, from [7], if (U, ϕ) is a minimal pair then we may choose a cancellation free pair (W, ψ) such that $\psi(W)$ is conjugate to $\phi(U)$ in $F(X)$. Then (W, ψ) is necessarily minimal. Hence $\mathcal{F}$ contains a cancellation free minimal pair. Moreover, if (W, ψ) is minimal and cancellation free then, for each element E of $\text{supp}(W)$, it follows that $\psi(E)$ is H-minimal.

Now, let (W, θ) be a minimal cancellation free pair in $\mathcal{F}$. If p is a geodesic path between 2 vertices of the Cayley graph $\Gamma_X(H)$ and p has label α then, by abuse of notation, we refer to the path p as α. In particular, for $(W, \theta) \in \mathcal{F}$, if $W = A_1 \cdots A_n$, where $A_i \in \mathcal{A}^{\pm 1}$, then we consider $\theta(W) = \theta(A_1) \cdots \theta(A_n)$ as the concatenation of paths $\theta(A_i)$ in the Cayley graph $\Gamma_X(H)$. This path contains a geodesic subpath with label $\theta(A_i)$, which we refer to as the path $\theta(A_i)$. On the other hand there is a geodesic path in $\Gamma_X(H)$ from 1 to $\theta(W)$, with label an H-minimal word F such that $F =_H \theta(W)$. We call such a path a *geodesic for* (W, θ) and refer to it as F.

Now suppose that, for each letter E of the Wicks form W, we have $|\theta(E)| \le L = 12l + M + 4$. In this case, in the light of Lemma 2.10, condition 1 of the appropriate Theorem is satisfied, apart from the bound on the length of R, which is deferred to Section 4.4. Therefore, from now on, we shall assume that there is at least one letter of W which is labelled by a word of length greater than L in $F(X)$. (This implies that there are two since each letter appears twice.)

For a pair $(W, \theta) \in \mathcal{F}$ and $A \in \text{supp}(W)$, we say that the occurrences of A and A^{-1} in W are *long edges of* (W, θ) (or just W) if $|\theta(A)| > L$, as a word in $F(X)$. All other letters of W are called *short edges*. For convenience in the proof, we may, without loss of generality, assume that the last letter of W is a long edge.

We state a number of preliminary lemmas, which we need later. The first is a version of Lemma 14 of [22], and establishes bounded length properties of subwords of $\theta(A)$, where A is any letter in $\text{supp}(W)$. Let Γ_W be the genus n graph associated to W.

Lemma 4.1. *Let (W, θ) be a minimal cancellation free pair in $\mathcal{F}$, let $A \in \operatorname{supp}(W)$, with signature $\sigma(A) = (\varepsilon, \delta)$, and let $E_1, \dots, E_r, A^\varepsilon, E_{r+1}, \dots, E_s, A^\delta, E_{s+1}, \dots, E_t$ be the cyclic sequence of letters in the Eulerian circuit W in Γ_W. Let $\theta(A^\varepsilon) = a_1 a_2$ and $\theta(A^\delta) = a'_1 a'_2$, where a_1, a'_1, a_2 and a'_2 are subwords of the H-minimal words $\theta(A^\varepsilon)$ and $\theta(A^\delta)$. Similarly, let $\theta(E_i) = e_{i1} e_{i2}$, for $1 \le i \le t$. Then*

(i) $|a_1| \le |e_{i2}\theta(E_{i+1}\cdots E_r)a_1|_H$, *for* $1 \le i \le r$;

(ii) $|a_2| \le |a_2\theta(E_{r+1}\cdots E_{j-1})e_{j1}|_H$, *for* $r+1 \le j \le s$;

(iii) $|a_1| \le |a_2\theta(E_{r+1}\cdots E_s A^\delta E_{s+1}\cdots E_{k-1})e_{k1}|_H$, *for* $s+1 \le k \le t$;

(iv) if $\delta = \varepsilon$ *and* $|a_1| \ge |a'_1|$ *then* $|a_2| \le |a_2\theta(E_{r+1}\cdots E_s)a'_1|_H$ *and*

(v) if $\delta = \varepsilon$ *and* $|a_2| \ge |a'_2|$ *then* $|a_1| \le |a_2\theta(E_{r+1}\cdots E_s)a'_1|_H$.

Proof. Without loss of generality we may assume that $\varepsilon = 1$ so $\sigma(A) = (1, \delta)$. First we shall prove statement (ii). As in Section 2.3 consider a disk D with its boundary divided into $t+2$ segments, labelled by the word $W = E_1 \cdots E_t$. Suppose that $E_j = X^\gamma$, for some $X \in \mathcal{A}$ and $\gamma = \pm 1$. Then there is (unique) $i \ne j$ such that $E_i = X^{\pm 1}$; say $E_i = E_j^\xi$, where $\xi = \pm 1$. Bisect E_j into two new edges, the first new edge denoted E_{j1} and the second E_{j2}, where E_{j1}, E_{j2} are elements of $\mathcal{A}^{\pm 1}$ not occurring in W. Also replace the edge E_i by the two edge path $E_{i1}E_{i2}$, where $E_{i1} = E_{j1}$ and $E_{i2} = E_{j2}$, if $\xi = 1$, and $E_{i1} = E_{j2}^{-1}$ and $E_{i2} = E_{j1}^{-1}$, if $\xi = -1$. That is, $E_{i1} = E^\xi_{j\mu(\xi)}$ and $E_{i2} = E^\xi_{j\nu(\xi)}$.

Now consider a properly embedded arc in D, with end points $\iota(A)$ and $\tau(E_{j1})$. Direct this arc from $\iota(A)$ to $\tau(E_{j1})$ and label it A'. (See Figure 4.1a.) Cut the disk D along A' to give two disks D_1 and D_2. Identify the segments of the boundaries of D_1 and D_2 labelled A to form a new disk D' with boundary labelled with the quadratic word



(a) (b) $\delta = -1$ (c) $\delta = 1$

Figure 4.1

$$W' = E_1 \cdots E_r A' E_{j2} \cdots E_s (E_{r+1} \cdots E_{j1} A'^{-1})^{-\delta} E_{s+1} \cdots E_t,$$

as shown in Figures 4.1b and 4.1c, for the cases $\delta = -1$ and $\delta = 1$, respectively. By considering links of vertices we see that W' is irredundant unless it contains, as a subword, two occurrences of either $(E_sE_{r+1})^{\pm 1}$, in the case $\delta = -1$; or $(E_{r+1}^{-1}E_{s+1})^{\pm 1}$, in the case $\delta = 1$. If this is the case, say $E_aE_b = E_{r+1}^{-1}E_s^{-1}$, then we replace the subword E_sE_{r+1} by a new letter E_u and the subword E_aE_b by E_u^{-1}. All other possibilities are dealt with similarly. Thus we may assume that W' is an irredundant quadratic word. As the surface Σ_W, obtained by identifying edges of D, according to their labels, is the same as the surface $\Sigma_{W'}$, the quadratic word W' is a genus n Wicks form. Let $\mathcal{L} = \operatorname{supp}(W')$ and let t_1 be an H-minimal word in $F(X)$, such that $t_1 =_H a_2\theta(E_{r+1}\cdots E_{j-1})e_{j1}$. Define a homomorphism $\psi : F(\mathcal{L}) \to F(X)$ in the following way.

$$\psi(E) = \begin{cases} a_1t_1 & \text{if } E = A' \\ e_{j1} & \text{if } E = E_{j1} \\ e_{j2} & \text{if } E = E_{j2} \\ e^{\xi}_{j\mu(\xi)} & \text{if } E = E_{i1} \\ e^{\xi}_{j\nu(\xi)} & \text{if } E = E_{i2} \\ \theta(E) & \text{otherwise.} \end{cases},$$

with the obvious adjustments if the word W' had to be modified to make it irredundant. Let

$$\begin{aligned} w_0 &= \theta(E_1\cdots E_r) = \psi(E_1\cdots E_r), \\ w_1 &= \theta(E_{r+1}\cdots E_{j-1})e_{j1} = \psi(E_{r+1}\cdots E_{j-1}E_{j1}), \\ w_2 &= e_{j2}\theta(E_{j+1}\cdots E_s) = \psi(E_{j2}E_{j+1}\cdots E_s) \text{ and} \\ w_3 &= \theta(E_{s+1}\cdots E_t) = \psi(E_{s+1}\cdots E_t). \end{aligned}$$

Then, writing E_i for $E_{i1}E_{i2}$, as before, so $\psi(E_i) = (e_{j1}e_{j2})^{\gamma\xi} = e_j^{\gamma\xi}$, we have

$$\begin{aligned} \psi(W') &=_H \psi(E_1\cdots E_rA'E_{j2}\cdots E_s(E_{r+1}\cdots E_{j1}A'^{-1})^{-\delta}E_{s+1}\cdots E_t) \\ &=_H w_0a_1t_1w_2(w_1t_1^{-1}a_1^{-1})^{-\delta}w_3 \end{aligned}$$

and substituting $t_1 =_H a_2w_1$ in this expression gives

$$\psi(W') =_H w_0a_1a_2w_1w_2(a_2^{-1}a_1^{-1})^{-\delta}w_3 =_H \theta(W).$$

This implies that $\psi(W')$ is conjugate to h in H. Thus (W', ψ) is an element of $\mathcal{F}$. Now since $|\theta(W)|$ was chosen to be minimal over all pairs in $\mathcal{F}$, it follows that

$$|\psi(W')| \;\geq\; |\theta(W)|. \tag{4.1}$$

Also, as θ is cancellation free on W, we have

$$\begin{aligned} |\theta(W)| &= |w_0| + |w_1| + |w_2| + |w_3| + 2|a_1| + 2|a_2| \\ &= (|w_0| + |w_1| + |w_2| + |w_3| + 2|a_1| + 2|t_1|) + 2|a_2| - 2|t_1| \\ &\geq |\psi(W')| + 2|a_2| - 2|t_1|. \end{aligned}$$

That is

$$|\theta(W)| \geq |\psi(W')| + 2|a_2| - 2|t_1|. \tag{4.2}$$

Inequalities (4.1) and (4.2) imply

$$|a_2\theta(E_{r+1}\cdots E_{j-1})e_{j1}|_H = |t_1| \geq |a_2|, \tag{4.3}$$

as required. Hence (ii) holds.

The same argument, using $(W^{-1}, \theta) \in \mathcal{F}$ and the cyclic permutation of W^{-1} beginning with E_S^{-1}, can be used to show that (i) holds.

Next we consider case (iii). Again we label the disk D with the word W. This time, bisect the edge labelled E_k. The first half shall be labelled by E_{k1} and the second half labelled by E_{k2}, where E_{k1}, E_{k2} are elements of $\mathcal{A}^{\pm 1}$ not occurring in W. As before there is an edge $E_i = E_k^{\eta}$, $\eta = \pm 1$, and we replace E_i with $E_{i1}E_{i2}$, where $E_{i1} = E^{\eta}_{k\mu(\eta)}$ and $E_{i2} = E^{\eta}_{k\nu(\eta)}$. Cut along an arc A'' joining $\tau(A)$ to $\tau(E_{k1})$, and identify the two edges labelled A, as in Figure 4.2. As before, if $\iota(A)$ in Γ_W has degree 3, and $\delta = -1$ then we also replace both occurrences the subword E_rE_{s+1} (or its inverse) with a new edge $E_u \in \mathcal{A}$ (or its inverse). A similar modification to the edge path $E_rE_s^{-1}$ may be necessary when $\delta = 1$. The boundary of the new disk is now labelled by the irredundant quadratic word



(a) (b) $\delta = -1$ (c) $\delta = 1$

Figure 4.2

$$W'' = E_1 \cdots E_r(E_{s+1} \cdots E_{k1}A''^{-1}E_{r+1} \cdots E_s)^{-\delta}A''E_{k2} \cdots E_t,$$

(replacing E_rE_{s+1} by E_u if necessary) which is a genus n Wicks form. We shall define a labelling function for W''. Let $\mathcal{K} = \mathrm{supp}(W'')$ and let t_2 be an H-minimal word in $F(X)$ such that

$$t_2 =_H a_2\theta(E_{r+1}\cdots E_s)(a_1a_2)^{\delta}\theta(E_{s+1}\cdots E_{k-1})e_{k1}. \tag{4.4}$$

Define a homomorphism $\phi : F(\mathcal{K}) \to F(X)$ by

$$\phi(E) = \begin{cases} a_2^{-1}t_2 & \text{if } E = A'' \\ e_{k1} & \text{if } E = E_{k1} \\ e_{k2} & \text{if } E = E_{k2} \\ e^{\eta}_{k\mu(\eta)} & \text{if } E = E_{i1} \\ e^{\eta}_{k\nu(\eta)} & \text{if } E = E_{i2} \\ \theta(E) & \text{otherwise.} \end{cases} .$$

As before we find that $\phi(W'') =_H \theta(W)$,

$$|\phi(W'')| \;\; \geq \;\; |\theta(W)| \tag{4.5}$$

and

$$|\theta(W)| \;\; \geq \;\; |\phi(W'')| - 2|t_2| + 2|a_1|. \tag{4.6}$$

Inequalities (4.5) and (4.6), imply that

$$|a_2\theta(E_{r+1}\cdots E_s)a_2^{-1}a_1^{-1}\theta(E_{s+1}\cdots E_{k-1})e_{k_1}|_H = |t_2| \geq |a_1|,$$

as required. Hence (iii) holds.

For (iv) we modify the arguments above as follows. As we are assuming $\varepsilon = 1$ we have $\delta = 1$. First consider the case where $i = 1$, and follow the proof of case (ii), but take $j = s$ and $E_{j1} = E_s$, $E_{j2} = 1$. Then

$$W' = E_1 \cdots E_r A'^2 E_s^{-1} \cdots E_{r+1}^{-1} E_{s+1} \cdots E_t.$$

Let E_0 be an element of $\mathcal{A}$ which does not belong to $\text{supp}(W')$ and replace W' with

$$W'_0 = E_1 \cdots E_r E_0 A'^2 E_0^{-1} E_s^{-1} \cdots E_{r+1}^{-1} E_{s+1} \cdots E_t.$$

The surface obtained from identifying boundary intervals of a disk labelled by W'_0 is again Σ_W, (as can be seen by considering the link of the vertex incident to A'. As $|a_1| \geq |a'_1|$, we have $a_1 = a'_1 a_3$, for some terminal subword a_3 of a_1. Define $t_1 =_H a_2\theta(E_{r+1}\cdots E_s)a'_1$, let $\mathcal{K}' = \mathcal{K} \cup \{E_0\}$ and define $\psi : F(\mathcal{K}') \to F(X)$ by

$$\psi(E) = \begin{cases} a_3 t_1 & \text{if } E = A' \\ a'_1 & \text{if } E = E_0 \\ \theta(E) & \text{otherwise} \end{cases} ,$$

w_0 and w_3 as before and

$$w_1 = \theta(E_{r+1}\cdots E_s) = \psi(E_{r+1}\cdots E_s).$$

Then $t_1 =_H a_2 w_1 a'_1$ so

$$\begin{aligned}\psi(W'_0) &=_H w_0 a'_1 a_3 t_1 a_3 t_1 a_1'^{-1} w_1^{-1} w_3\\ &=_H w_0 a_1 a_2 w_1 a_1 a_2 w_1 a'_1 a_1'^{-1} w_1^{-1} w_3\\ &=_H w_0 a_1 a_2 w_1 a_1 a_2 w_3\\ &=_H \theta(W).\end{aligned}$$

As before, W' is a genus n Wicks form so $|\theta(W)| \leq |\phi(W'_0)|$. Thus

$$|w_0| + |w_1| + |w_3| + 2|a_1| + 2|a_2| = \theta(W) \leq \phi(W'_0) \leq |w_0| + |w_1| + |w_3| + 2|a'_1| + 2|a_3| + 2|t_1|,$$

which implies that $|t_1| \geq |a_2|$, so (iv) holds.

To prove that (v) holds, consider again the word W'', but this time take $k = s + 1$, $E_{k1} = 1$ and $E_{k2} = E_{s+1}$. Then

$$W'' = E_1 \cdots E_r E_s^{-1} E_{r+1}^{-1} \cdots A''^2 E_{s+1} \cdots E_t.$$

Let E_0 be an element of $\mathcal{A}$ which does not belong to $\text{supp}(W'')$ and replace W'' with

$$W''_0 = E_1 \cdots E_r E_s^{-1} \cdots E_{r+1}^{-1} E_0^{-1} A''^2 E_0 E_{s+1} \cdots E_t.$$

The surface obtained from identifying boundary intervals of a disk labelled by W''_0 is again Σ_W. As $|a_2| \geq |a'_2|$, we have $a_2 = a_4 a'_2$, for some initial subword a_4 of a_2, so $a'_1 = a_1 a_4$. Again, let $t_1 =_H a_2 \theta(E_{r+1} \cdots E_s) a'_1$, and define

$$\phi(E) = \begin{cases} a_4^{-1} t_1 & \text{if } E = A'' \\ a'_2 & \text{if } E = E_0 \\ \theta(E) & \text{otherwise} \end{cases},$$

and w_0, w_1, w_3 as in case (iv). Then

$$\begin{aligned}\phi(W''_0) &=_H w_0 w_1^{-1} a_2'^{-1} (a_4^{-1} t_1)^2 a'_2 w_3\\ &= w_0 w_1^{-1} a_2'^{-1} (a_4^{-1} a_2 w_1 a'_1)^2 a'_2 w_3\\ &= w_0 a'_1 a_4^{-1} a_2 w_1 a'_1 a'_2 w_3\\ &= \theta(W).\end{aligned}$$

Again, W''_0 is a genus n Wicks form so $|\theta(W)| \leq |\phi(W''_0)|$. Thus

$$|w_0| + |w_1| + |w_3| + 2|a_1| + 2|a_2| = \theta(W) \leq \phi(W''_0) \leq |w_0| + |w_1| + |w_3| + 2|a'_2| + 2|a_4| + 2|t_1|,$$

and consequently $|t_1| \geq |a_1|$, so (v) holds. □

Next we record as corollaries some straightforward consequences of Lemma 4.1.

Corollary 4.2. *Let (W, θ) be a cancellation free pair in $\mathcal{F}$ and let F be a geodesic for (W, θ). Let A be a long edge of W, with $\sigma(A) = (\varepsilon, \delta)$. Let $\alpha = \theta(A^\varepsilon)$ and $\alpha' = \theta(A^\delta)$, let x be a vertex of α and k be a positive integer such that $k < d(\iota(\alpha), x) < |\alpha| - k$. Then the following hold.*

(i) *Let y be a vertex on $(\theta(W) \cup F) - \alpha$ such that $d(x, y) \leq k$. Then y lies on $\alpha' \cup F$ and if $\delta = \varepsilon$ then y lies on F.*

(ii) *If $k > l$ then there exists a vertex y such that y lies on $\alpha' \cup F$ and $d(x, y) \leq l$. If $\delta = \varepsilon$ then y lies on F.*

Proof. Statement (i) follows from Lemma 4.1. For statement (ii) note that by Lemma 2.3.(i), there exists a vertex y on $\theta(W) - \alpha$ such that $d(x, y) \leq l$. As $k > l$ the statement now follows from (i). □

Corollary 4.3. *Let (W, θ), F, A, α and α' be as in Corollary 4.2. Let x_1 and x_2 be any vertices on α and α', respectively, such that $d(x_1, x_2) \leq k$ for some constant k. If x_3 is a vertex on α' such that $d(\iota(\alpha'), x_3) = d(\tau(\alpha), x_1)$ then $d(x_2, x_3) \leq k$ and $d(x_1, x_3) \leq 2k$.*

Proof. The proof falls into the following two cases.

(a) $d(\iota(\alpha'), x_2) \leq d(\iota(\alpha'), x_3) = d(\tau(\alpha), x_1)$.

(b) $d(\iota(\alpha'), x_2) > d(\iota(\alpha'), x_3) = d(\tau(\alpha), x_1)$.

(a) By Lemma 4.1 $d(\tau(\alpha), x_1) \leq d(x_1, \iota(\alpha'))$ and, from the hypothesis and the triangle inequality, it follows that

$$d(\tau(\alpha), x_1) \leq d(x_1, \iota(\alpha')) \leq k + d(\iota(\alpha'), x_2). \tag{4.7}$$

Since α is a geodesic path we have

$$d(\tau(\alpha), x_1) = d(\iota(\alpha'), x_3) = d(\iota(\alpha'), x_2) + d(x_2, x_3). \tag{4.8}$$

It follows from equations (4.7) and (4.8) that $d(x_2, x_3) \leq k$.

(b) By Lemma 4.1, $d(\iota(\alpha'), x_2) \leq d(x_2, \tau(\alpha))$ and, from the hypothesis and the triangle inequality, it follows that

$$d(\iota(\alpha'), x_2) \leq d(x_2, \tau(\alpha)) \leq k + d(\tau(\alpha), x_1). \tag{4.9}$$

Since α is a geodesic path we have

$$d(\tau(\alpha), x_1) = d(\iota(\alpha'), x_3) = d(\iota(\alpha'), x_2) - d(x_2, x_3). \tag{4.10}$$

It follows from equations (4.9) and (4.10) that $d(x_2, x_3) \leq k$. Hence the lemma holds. □

Corollary 4.4. *Let (W, θ), F, A, α and α' be as in Corollary 4.2. Assume in addition that $o(A) = 1$, so $\sigma(A) = (\varepsilon, -\varepsilon)$ and $\alpha' = \alpha^{-1}$. For $i = 1, 2$, let u_i be vertices of α and v_i be vertices of α^{-1} such that*

- $d(\tau(\alpha), u_1) \geq d(\tau(\alpha), u_2)$ *and*
- *for some constant k, $d(u_i, v_i) \leq k$, $i = 1, 2$.*

Let α_1 *be the subpath* $[u_1, u_2]$ *of* α. *(See Figure 4.3.) Then* $|\alpha_1| \leq 2k + M + 1$.

Proof. Let w_i be the vertex of α^{-1} such that $d(\tau(\alpha^{-1}), w_i) = d(\iota(\alpha), u_i)$, $i = 1, 2$. Let s_i be a geodesic from u_i to w_i, for $i = 1, 2$. From Corollary 4.3, we have $|s_i| \leq 2k$. The subpath $[w_1, w_2]$ of α^{-1} is α_1^{-1}, so from Lemma 2.2, we have

$$|\alpha_1| \leq \frac{1}{2}(|s_1| + |s_2|) + M + 1 \leq 2k + M + 1.$$

□



Figure 4.3

We shall use Corollaries 4.2 and 4.4 with Lemma 2.3 to factor F into a product of subwords, each of which corresponds to a unique long edge of W. To this end let $W = E_1 \cdots E_m$, where $E_i = A_i^{\varepsilon_i}$, with $A_i \in \mathcal{A}$ and $\varepsilon_i \in \{\pm 1\}$. Let U be the word obtained from W by deleting E_i, whenever E_i is a short edge. Then U is a non-trivial quadratic word over $\mathcal{A}$, of genus at most n, and length f at most $K(n)$, which we call the *long factorisation* of W (with respect to θ). Write $U = U_1 \cdots U_f$, where $U_i \in \{E_1, \dots, E_m\}$, and U_i is a long edge of W. Then, as W ends with a long edge, we can write $W = S_1 U_1 S_2 \cdots S_f U_f$, where each S_i is a subword of W consisting only of short edges (and may be the empty word). By the usual abuse of notation, let $\theta(U_i)$ and $\theta(S_i)$ denote subpaths in $\Gamma_X(H)$ of the path labelled $\theta(W)$.

Lemma 4.5. *In the above notation, let* $W = S_1 U_1 S_2 \cdots S_f U_f$, *where* $U = U_1 \cdots U_f$ *is the long factorisation of* W *and each* S_i *is a product of short edges. Then* F *has a partition* $F_1, \dots, F_f$ *such that*

1. *if* $o(U_i) = -1$ *then* $|F_i| \geq 8l + M + 3$ *and* $d(\tau(F_i), \tau(\theta(U_i)) \leq 3l + 1$*;*
2. *if* $o(U_i) = 1$ *then* $|F_i| \geq 6l + 2$ *and, for the unique* j *such that* $U_i = U_j^{-1}$, *we have, assuming* $i < j$, *that* $d(\tau(F_i), \tau(\theta(U_i)) \leq 5l + M + 3$ *and* $d(\tau(F_i), \tau(\theta(U_j)) \leq 3l + 1$,

for all i *such that* $1 \leq i \leq f$.

Proof. Let $A \in \operatorname{supp}(W)$ such that A is a long edge and assume that $W = W_0U_iW_0U_jW_1$, where $U_i = A^\varepsilon$ and $U_j = A^\delta$. Let $\alpha = \theta(A^\varepsilon)$ and $\beta = \theta(A^\delta)$. Then $|\alpha| = |\beta| > 12l + M + 4$ and factorising as $\alpha = \alpha_0\alpha_1\alpha_2$, where $|\alpha_0| = |\alpha_2| = 2l + 1$ we have $|\alpha_1| > 8l + M + 2$.

Now, as above, $W = E_1 \cdots E_m$, where $E_i = A_i^{\varepsilon_i}$, with $A_i \in \mathcal{A}$ and $\varepsilon_i \in \{\pm 1\}$. Apply Lemma 2.3 to the word $\theta(E_1)\cdots\theta(E_m)F^{-1}$, with $\gamma_0 = \theta(U_i)$. This implies there is a partition $\gamma^{(1)}, \dots, \gamma^{(r)}$ of α such that, for each $j \in \{1, \dots, r\}$, the interval $\gamma^{(j)}$ is a $\delta\lceil\log_2(m)\rceil$-fellow traveller with a geodesic γ', which is either an interval of a partition of $\theta(E_s)$, for some $E_s \neq U_i$, or of F^{-1}. As $\delta\lceil\log_2(m)\rceil \le l$, it follows that $\gamma^{(j)}$ and γ' are l-fellow travellers.

Assume that the partition of F^{-1} given by Lemma 2.3 is $F'_1, \dots, F'_t$. Suppose first that $o(A) = -1$ and $\sigma(A) = (\varepsilon, \varepsilon)$. Then, as $|a_0| = |a_2| = 2l + 1$, it follows from Corollary 4.2, that α_1 is a subinterval of $\gamma^{(b)}$, where $1 \le d \le r$ and $\gamma^{(b)}$ is an l-fellow traveller with F'_c, for some c, and that $|F'_c| \ge |a_1| \ge 8l + M + 3$. In this case, a similar argument shows that a subinterval $\gamma^{(b')}$ of β is an l-fellow traveller with $F'_{c'}$, for some $c' < c$, with $|F'_c| \ge 8l + M + 3$.

This leaves the case $o(A) = 1$ and $\sigma(A) = (\varepsilon, -\varepsilon)$ to consider. In this case, from Corollary 4.2, α_1 is a subinterval of $\gamma^{(b)}\gamma^{(b+1)}$, where γ^b is an l-fellow traveller with an interval F'_c of F'^{-1} and $\gamma^{(b+1)}$ is an l-fellow traveller with an interval $\gamma'^{(d)}$ of the partition $\gamma'^{(1)}, \dots, \gamma'^{(r')}$ of β given by Lemma 2.3. Thus $|\gamma^{(b)}| + |\gamma^{(b+1)}| = |\gamma^{(b)}\gamma^{(b+1)}| \ge |\alpha_1| \ge 8l + M + 3$. From Corollary 4.4, $|\gamma^{(b+1)}| \le 2l + M + 1$, so $|F'_c| = |\gamma^{(b)}| \ge 6l + 2$. Moreover, in this case the interval $\gamma'^{(d+1)}$ is an l-fellow traveller with an interval $F'_{c'}$ of the partition of F'^{-1}, with $c' < c$, and again $|F'_{c'}| \ge 6l + 2$.

Recalling that $\alpha = \theta(U_i)$ and $\beta = \theta(U_j)$, in both cases we denote the initial vertex $\iota(F'_c)$ of F'_c by v_i and the initial vertex $\iota(F'_{c'})$ of $F'_{c'}$ of by v_j. By construction, if $o(A) = -1$, then $d(v_i, \tau(\theta(U_i))) \le 3l+1$ and if $o(A) = 1$, then $d(v_i, \tau(\theta(U_i))) \le 3l+1+2l+M+1 = 5l+M+2$, while in both cases $d(v_j, \tau(\theta(U_j))) \le 3l + 1$.

Define v_i and v_j in this way for each long edge $A \in \operatorname{supp}(W)$. Setting $v_0 = \tau(F)$ results in $f+1$ points $v_0, v_1, \dots, v_f$ on F such that $d(v_0, v_i) > d(v_0, v_{i+1})$ and, with $F_{f-i} = [v_{i+1}, v_i]$, we have a partition $F_1, \dots, F_f$ of F, where $|F_i| \ge 6l + 2$. $\square$

Definition 4.6. *With the notation of the paragraph above Lemma 4.5, let* $W = S_1U_1 \cdots S_fU_f$, *let* $U = U_1 \cdots U_f$ *be the long factorisation of* W, *let* $F = F_1 \cdots F_f$ *be the factorisation of* F *found in Lemma 4.5 and let* $v(U_i) = \tau(F_i)$. *For* $i = 1, \dots, l$, *let* $a(U_i)$ *be the label of a geodesic path from* $\tau(\theta(U_i))$ *to* $v(U_i)$, *let* $s(U_i) = \theta(S_i)$ *(and let* $a(U_0) = 1$ *and* $S_0 = 1$*) and let* $\alpha(U_i) = F_i$. *For later reference let* $S(U_i) = S_i$, *for all* i. *Let* $A \in \operatorname{supp}(U)$.

(i) If $o(A) = 1$, *let* i, j *be such that* $U_i = A$ *and* $U_j = A^{-1}$.

(ii) If $o(A) = -1$ *and* $\sigma(A) = (\varepsilon, \varepsilon)$, *let* i, j *be such that* $U_i = U_j = A^\varepsilon$, $i < j$.

Define

$$v_1(A) = v(U_i), \quad a_1(A) = a(U_i), \quad b_1(A) = a(U_{i-1}), \quad S_1(A) = S_i, \quad s_1(A) = s(U_i)$$

$$\textit{and } \alpha_1(A) = \begin{cases} \alpha(U_i), & \textit{if } o(A) = 1, \\ \alpha(U_i)^{\varepsilon}, & \textit{if } o(A) = -1; \end{cases}$$

$$v_2(A) = v(U_j), \quad a_2(A) = a(U_j), \quad b_2(A) = a(U_{j-1}), \quad S_2(A) = S_j, \quad s_2(A) = s(U_j)$$

$$\textit{and } \alpha_2(A) = \begin{cases} \alpha(U_j)^{-1}, & \textit{if } o(A) = 1, \\ \alpha(U_j)^{\varepsilon}, & \textit{if } o(A) = -1. \end{cases}$$

(See Figure 4.4.)

Finally define H-minimal words

$$z_1(A) =_H \begin{cases} b_2(A)^{-1} s_2(A) a_1(A), & \textit{if } o(A) = 1, \\ a_2(A)^{-1} a_1(A), & \textit{if } o(A) = -1, \end{cases} \quad \textit{and}$$

$$z_2(A) =_H \begin{cases} b_1(A)^{-1} s_1(A) a_2(A), & \textit{if } o(A) = 1, \\ b_1(A)^{-1} s_1(A) s_2(A)^{-1} b_2(A), & \textit{if } o(A) = -1. \end{cases}$$



(a) $o(A) = 1$, $A = U_i$, $A^{-1} = U_j$, $i < j$ (b) $o(A) = 1$, $A = U_i$, $A^{-1} = U_j$, $i > j$



(c) $o(A) = -1$, $A = U_i = U_j$ (d) $o(A) = -1$, $A^{-1} = U_i = U_j$

Figure 4.4

Example 4.7. Suppose that $W = ABC^{-1}A^{-1}B^{-1}C^{-1}$ where A and C are long edges and B is a short edge. Then $U = AC^{-1}A^{-1}C^{-1}$ and we have the paths in $\Gamma_X(H)$ shown in Figure 4.5. Here $|a_1(A)|$, $|a_2(A)|$ and $|a_1(C)|$ are at most $5l + M + 3$ and $|a_2(C)| = 0$. (As in our standing assumption, the last letter of W is long). In this case $S_0 = S_2 = S_4 = 1$,



Figure 4.5

$S_1 = B$ and $S_3 = B^{-1}$, so $b_1(A) = 1$, $b_2(A) = a_1(C)$, $b_1(C) = a_1(A)$ and $b_2(C) = a_2(A)$, $s_i(A) = 1$, $s_1(C) = \theta(B)$ and $s_2(C) = \theta(B)^{-1}$. As $o(A) = 1$ and $o(C) = -1$, $z_i(A)$ and $z_i(C)$ are H-minimal elements of $F(X)$ such that

$z_1(A) = (a_1(C))^{-1}a_1(A)$, $\quad z_2(A) = a_2(A)$, $\quad z_1(C) = a_1(C)$ and $z_2(C) = (a_1(A))^{-1}\theta(B)^2 a_2(A)$.

We shall need the following lemma in the next subsection.

Lemma 4.8. *Let A be a long edge of W. Then*

(i) $z_2(A)\alpha_2(A)z_1(A)\alpha_1(A)^{-1} =_H 1$, if $o(A) = 1$, and

(ii) $z_2(A)\alpha_2(A)^{\varepsilon}z_1(A)\alpha_1(A)^{-\varepsilon} =_H 1$, if $o(A) = -1$ and $\sigma(A) = (\varepsilon, \varepsilon)$.

Proof. By definition, for $1 \leq i \leq l$ the Cayley graph $\Gamma_X(H)$ contains a closed path, based at $v(U_{i-1})$ on F, consisting of the concatenation of paths labelled $\alpha(U_i)$, $a(U_i)^{-1}$, $\theta(U_i)^{-1}$, $\theta(S_{i-1})^{-1}$ and $a(U_{i-1})$. (See Figure 4.6). Thus, for $A \in \mathrm{supp}(U)$ with $o(A) = 1$, we have both

$$\alpha_1(A)a_1(A)^{-1}\theta(A)^{-1}s_1(A)^{-1}b_1(A) =_H 1 \text{ and } \alpha_2(A)^{-1}a_2(A)^{-1}\theta(A)s_2(A)^{-1}b_2(A) =_H 1.$$

Similarly, for $A \in \mathrm{supp}(U)$ with $o(A) = -1$, we have

$$\alpha_k(A)^{\varepsilon}a_k(A)^{-1}\theta(A)^{-\varepsilon}s_k(A)^{-1}b_k(A) =_H 1, \text{ for } k = 1, 2.$$

The result follows from these two facts and the definition of $z_i(A)$. $\square$



(a) $o(A) = 1$, $A = U_i$, $A^{-1} = U_j$

(b) $o(A) = -1$, $\sigma(A) = (\varepsilon, \varepsilon)$

Figure 4.6

4.2 Construction of an extension of U

Write W around the boundary of a disk (i.e. divide the boundary up into $|W|$ segments, assigning a letter to each) and identify the long edges, respecting orientation. This results in a surface S of genus $k \leq n$ with Q holes. The boundary of the disk becomes a graph on this surface, which we denote by Γ_S. Thus Γ_S consists of short edges, all of which are written around the boundary components of S, and long edges, all of which are properly embedded on S.

Let C be the set of cyclic words, over $\mathcal{A}$, labelling the Q boundary components of S. Define $\sim$ to be the equivalence relation on C generated by the relation which consists of all pairs (U, V) of words in C such that $\mathrm{supp}_{F(\mathcal{A})}(U) \cap \mathrm{supp}_{F(\mathcal{A})}(V) \neq \emptyset$. Let $B_1, \ldots, B_p$ be the equivalence classes of $\sim$. Then each set B_i consists of a t_i-tuple of cyclic words $(W^i_1, \ldots, W^i_{t_i})$, where W^i_j is a cycle forming a boundary component of S, and by definition of B_i, this t_i-tuple is quadratic.

Definition 4.9. *The* S-genus *of* B_i *is* $g_i = \mathrm{genus}_H(\theta(W^i_1), \ldots, \theta(W^i_{t_i})) + t_i - 1$, *for* $i = 1, \ldots, p$.

Lemma 4.10. *Let* $g = \sum_{i=1}^{p} g_i$, *where* g_i *is the* S*-genus of* B_i, *and let* $\mathrm{genus}(S) = k$. *For each equivalence class* B_i *consisting of a* t_i*-tuple* $(W^i_1, \ldots, W^i_{t_i})$ *we have*

$$\mathrm{genus}_H(\theta(W^i_1), \ldots, \theta(W^i_{t_i})) = \mathrm{genus}_{F(\mathcal{A})}(W^i_1, \ldots, W^i_{t_i}), \quad \textit{for } i = 1, \ldots, p.$$

It follows that $g + k = n$.

Proof. Let $h_i = \mathrm{genus}_{F(\mathcal{A})}(W^i_1, \ldots, W^i_{t_i}) + t_i - 1$. Then $h_i \geq g_i$, for all i. Assume that $h_j > g_j$, for some $1 \leq j \leq p$. Now, since W has genus n in $F(\mathcal{A})$, if we identify all the short edges on the genus k surface S, respecting orientation, we obtain the closed compact surface Σ_W of genus n. On the other hand we may construct a surface homeomorphic to

Σ_W as follows. For each B_i there exists a van Kampen diagram K_i, over $F(\mathcal{A})$, on a surface Σ_i of genus $h_i - t_i + 1$, with t_i boundary components, $W^j_1, \dots, W^j_{t_i}$. Identifying boundary components of Σ_i to the t_i boundary components of S, with the same boundary cycles, for all i, we obtain a closed surface of genus $k + \sum_{i=1}^p [(h_i - t_i + 1) + t_i - 1] = k + \sum_{i=1}^p h_i$. By construction this surface is homeomorphic to Σ_W.

Therefore,

$$\begin{aligned} n &= k + \sum_{i=1}^{p} h_i \\ &> k + \sum_{i=1}^{p} g_i. \end{aligned}$$

But this implies that $\text{genus}_H(h) = \text{genus}_H(\theta(W)) \le k + \sum_{i=1}^p g_i < n$, a contradiction. Thus $g_i = h_i$, for all $i = 1, \dots, p$, and $n = k + \sum g_i = k + g$. Hence the lemma holds. $\square$

Consider once more the surface S with Q holes with the embedded graph Γ_S consisting of long and short edges. If we paste a disk onto each of the Q boundary components and contract the cyclic word of short edges to a point, we obtain a graph, consisting of long edges only, on a closed compact surface of genus $k = \text{genus}(S)$. The quadratic word associated with this graph is the long factorisation U of W. As usual we denote this graph Γ_U. We shall construct an extension of U over H.

As above, let the long factorisation of W be $U = U_1 \cdots U_l$. As in Section 2.5, let U' the Hamiltonian cycle associated to U: that is the word obtained from U by replacing each occurrence U_i by U_{i1} or U_{i2}, as described in Section 2.5. Thus the word $U = AC^{-1}A^{-1}C^{-1}$ in Example 4.7 is replaced by $U' = A_1C_1^{-1}A_2^{-1}C_2^{-1}$. The graph obtained from Γ_U by replacing the directed edge U_i with edges U_{i1} and U_{i2} is called Γ'_U, as before. Now define a labelling function ψ on the set $\{A_1, A_2 : A \in \text{supp}(U)\}$, and by extension on the free group these letters generate, by

$$\psi(A_1) = \alpha_1(A) \text{ and } \psi(A_2) = \alpha_2(A).$$

By definition of α_1 and α_2 we have then

$$F = \psi(U'). \tag{4.11}$$

For instance, in Example 4.7 we have $\psi(U') = \alpha_1\gamma_1^{-1}\alpha_2^{-1}\gamma_2^{-1} = F$.

Let v^i_j be the vertex of Γ_U obtained by collapsing the boundary component W^i_j of S to a point. Let $B'_i = \{v^i_1, \dots, v^j_{t_i}\}$. We shall construct a genus g_i extension of Γ_U on B'_i, for appropriate g_i, for all i (see Definition 2.17). Each internal vertex of Γ_S becomes a vertex of Γ_U and we shall construct a genus 0 extension on each such vertex.

Let v be a vertex of Γ_U of degree d and assume that, after suitable relabelling of edges, the vertex sequence of $\text{lk}(v)$ is $E_1^{\varepsilon_1}, \dots, E_d^{\varepsilon_d}$, and the incidence sequence of v is $O_1, \dots, O_d$. Then the cyclic word U contains the subwords $(E_q^{\varepsilon_q}E_{q+1}^{\varepsilon_{q+1}})^{O_q}$, for $q = 1, \dots, d$, as in Figure 2.7a (with "e" replaced throughout by "E".)

Now extend each vertex v of Γ_U by a cycle graph C_v, as in Step 2 of the definition of an extension in Section 2.5. As in the definition, if the cycle $C_v = c_1 \cdots c_d$, where d is the degree of v and $\iota(c_q) = v_q$, $\tau(c_q) = v_{q+1}$, then using the notation above, we shall have $v_q = \tau((E^q_{\nu_q})^{\varepsilon_q}) = \tau((E^{q-1}_{\mu_{q-1}})^{\varepsilon_{q-1}})$, for $q = 1, \ldots d$ (superscripts modulo d), as in Figure 2.8c. The resulting graph is Γ''_U.

To complete Step 3 of the construction of the extension we must extend the labelling function ψ, defined above on A_1 and A_2, for $A \in \operatorname{supp}(U)$, to the edges of the cycles added to form Γ''_U. For the cycle C_v above we define, for $q = 1, \ldots, d$,

$$\psi(c_q) = z_{\mu_q}(E_q)^{O_q}, \text{ if } o(E_q) = 1, \tag{4.12}$$

and

$$\psi(c_q) = z_{\mu(O_q)}(E_q)^{\gamma_q}, \text{ if } o(E_q) = -1, \tag{4.13}$$

where

$$\gamma_q = O_q \nu^{-1}(l_q) = \begin{cases} -O_q, & \text{if } l_q = 1 \\ O_q, & \text{if } l_q = 2. \end{cases}.$$

Lemma 4.11. *The pair (Γ''_U, ψ) is an extension of U over H.*

Proof. We must verify that (a), (b) and (c), of Step 3 of the definition of extension, hold. By definition of ψ, (a) holds. To see that (b) holds, consider a long edge A of Γ_U and suppose that A is replaced by the pair (A_1, A_2), in forming Γ''_U. We have defined $\psi(A_i) = \alpha_i(A)$, for $i = 1, 2$. Assume that $\iota(A) = u$ and $\tau(A) = v$. Without loss of generality we may assume that the links of u and v have vertex sequences $E_1^{\varepsilon_1}, E_2^{\varepsilon_2}, \ldots, E_d^{\varepsilon_d}$ and $G_1^{\delta_1}, G_2^{\delta_2}, \ldots, G_f^{\delta_f}$, with $E_1^{\varepsilon_1} = A^{-1}$ and $G_1^{\delta_1} = A$. Write $O_q = O_q(u)$ and $O'_q = O_q(v)$, and similarly write r_q, l_q, μ_q and ν_q, for $r_q(u)$, $l_q(u)$, $\mu_q(u)$ and $\nu_q(u)$; and r'_q, l'_q, μ'_q and ν'_q, for $r_q(v)$, $l_q(v)$, $\mu_q(v)$ and $\nu_q(v)$. We may assume (by reversing orientation of all vertices if necessary) that $O_1 = 1$. From Lemma 2.15, the cyclic word U' contains

$$A_{r_1}^{-1} E_{2,l_2}^{-\varepsilon_2} \text{ and } (E_{d,r_d}^{\varepsilon_d} A_{l_1})^{O_d} \tag{4.14}$$

and

$$(A_{r'_1} G_{2,l'_2}^{-\delta_2})^{O'_1} \text{ and } (G_{f,r'_f}^{\delta_f} A_{l'_1}^{-1})^{O'_f}. \tag{4.15}$$

Assume further that u and v are extended by cycles C_u and C_v, where $C_u = c_1 \cdots c_d$ and $C_v = c'_1 \cdots c'_f$, with

$$\iota(c_q) = v_q, \tau(c_q) = v_{q+1}, \iota(c'_q) = v'_q \text{ and } \tau(c'_{q+1}) = v'_{q+1}.$$

Then

$$\iota(A_{l_1}) = v_1 = \iota(c_1), \iota(A_{r_1}) = v_2 = \tau(c_1), \tau(A_{l'_1}) = v'_1 = \iota(c'_1) \text{ and } \tau(A_{r'_1}) = v'_2 = \tau(c'_1).$$

Consider first the case $o(A) = 1$. As $\varepsilon_1 = -1$ and $\delta_1 = 1$, we have $\mu_1 = 2$, $\nu_1 = 1$, $\mu'_1 = 1$ and $\nu'_1 = 2$. Then

$$\psi(c_1) = z_{\mu_1}(A) = z_2(A) \tag{4.16}$$

and

$$l_1 = \nu_1 = 1, r_1 = \mu_1 = 2, \iota(A_1) = \iota(c_1) \text{ and } \iota(A_2) = \tau(c_1).$$

If $O'_f = 1$ then

$$\psi(c'_1) = z_{\mu'_1}(A) = z_1(A) \tag{4.17}$$

and

$$l'_1 = \nu'_1 = 2, r'_1 = \mu'_1 = 1, \tau(A_2) = \iota(c'_1) \text{ and } \tau(A_1) = \tau(c'_1).$$

We therefore have a closed path $c_1 A_2 c'_1 A_1^{-1}$ and, from Lemma 4.8 (i),

$$\psi(c_1)\psi(A_2)\psi(c'_1)\psi(A_1^{-1}) = z_2(A)\alpha_2(A)z_1(A)\alpha_1(A)^{-1} =_H 1,$$

so (b) holds in this case. If $O'_f = -1$ then

$$\psi(c'_1) = z_{\mu'_1}(A)^{-1} = z_1(A)^{-1} \tag{4.18}$$

and a similar argument shows that we have a closed path $c_1 A_2 {c'_1}^{-1} A_1^{-1}$; so (b) also holds in this case.

Now suppose that $o(A) = -1$. Then, from (4.14), U' contains A^{-1}, so $\sigma(A) = (-1, -1)$. From (4.15), then $O'_1 = -1$. Hence

$$\psi(c_1) = z_1(A)^{\nu^{-1}(l_1)} \text{ and } \psi(c'_1) = z_2(A)^{-\nu^{-1}(l'_1)}. \tag{4.19}$$

If $r_1 = l'_1$ and $l_1 = r'_1$ then

$$\psi(c'_1) = z_2(A)^{-\nu^{-1}(r_1)} = z_2(A)^{\nu^{-1}(l_1)},$$

and

$$\tau(A_{r_1}) = \tau(A_{l'_1}) = \iota(c'_1) \text{ and } \tau(A_{l_1}) = \tau(A_{r'_1}) = \tau(c'_1),$$

so we have a closed path $c_1 A_{r_1} c'_1 A_{l_1}^{-1}$ and

$$\psi(c_1)\psi(A_{r_1})\psi(c'_1)\psi(A_{l_1}^{-1}) = z_1(A)^{\nu^{-1}(l_1)}\alpha_{r_1}(A)z_2(A)^{\nu^{-1}(l_1)}\alpha_{l_1}(A)^{-1}.$$

Thus, as $\varepsilon = -1$, if $l_1 = 1$ we have

$$\psi(c_1)\psi(A_{r_1})\psi(c'_1)\psi(A_{l_1}^{-1}) = z_1(A)^{-1}\alpha_2(A)^{-\varepsilon}z_2(A)^{-1}\alpha_1(A)^{\varepsilon}$$

and if $l_1 = 2$ then

$$\psi(c_1)\psi(A_{r_1})\psi(c'_1)\psi(A_{l_1}^{-1}) = z_1(A)^{-1}\alpha_1(A)^{-\varepsilon}z_2(A)\alpha_1(A)^{\varepsilon}$$

and in both cases (b) follows from Lemma 4.8 (ii). The case $r_1 = r'_1$ and $l_1 = l'_1$ follows in the same way, so (b) holds in all cases.

As $\psi(U') = F$, (c) holds. Therefore we have an extension of U over H, as required. □

4.3 Length and genus of (Γ''_U, ψ)

The next two Lemmas prepare the ground for the proof of Lemma 4.14.

Lemma 4.12. *Let v^i_j be the vertex of Γ_U obtained by collapsing the boundary component W^i_j of S to a point. Let $E_1^{\varepsilon_1}, \dots, E_d^{\varepsilon_d}$ be the vertex sequence of $\mathrm{lk}(v^i_j)$, and $O_1, \dots, O_d$ the incidence sequence of v^i_j. Then $\theta(W^i_j)$ is conjugate to $u_1^{O_1} \cdots u_d^{O_d}$, where*

$$u_q = \begin{cases} s_{l_{q+1}}(E_{q+1}), & \textit{if } O_q = 1 \\ s_{r_q}(E_q), & \textit{if } O_q = -1. \end{cases}$$

Proof. By definition of O_q, the cyclic word U contains $(E_q^{\varepsilon_q} E_{q+1}^{-\varepsilon_{q+1}})^{O_q}$, and hence W contains $(E_q^{\varepsilon_q} S_q'^{O_q} E_{q+1}^{-\varepsilon_{q+1}})^{O_q}$, where

$$S'_q = \begin{cases} S(E_{q+1}^{-\varepsilon_{q+1}}), & \text{if } O_q = 1, \\ S(E_q^{-\varepsilon_q}), & \text{if } O_q = -1, \end{cases}$$

for $q = 1, \dots, d$. Moreover, from Lemma 2.15, the cyclic word U' contains $(E_{q,r_q}^{\varepsilon_q} E_{q+1,l_{q+1}}^{-\varepsilon_{q+1}})^{O_q}$, in place of $(E_q^{\varepsilon_q} E_{q+1}^{-\varepsilon_{q+1}})^{O_q}$. Hence, by considering all the possible values when $O_q = \pm 1$, $o(E_{q+1}) = \pm 1$ and $\varepsilon_{q+1} = \pm 1$, we see that if $O_q = 1$ then $S'_q = S_{l_{q+1}}(E_{q+1})$, while if $O_q = -1$ then $S'_q = S_{r_q}(E_q)$. ☐

In the following lemma we abbreviate the notation of Definition 4.6, writing a_{l_q}, b_{l_q}, s_{l_q}, a_{r_q}, b_{r_q} and s_{r_q} for $a_{l_q}(E_q)$, $b_{l_q}(E_q)$, $s_{l_q}(E_q)$, $a_{r_q}(E_q)$, $b_{r_q}(E_q)$ and $s_{r_q}(E_q)$, respectively.

Lemma 4.13. *Let v be a vertex of Γ_U with incidence sequence $O_1, \dots, O_d$ and link $E_1^{\varepsilon_1}, \dots, E_d^{\varepsilon_d}$. Let $C_v = c_1 \cdots c_d$ be the cycle extending v, let $V_q = (o(E_q), O_q; o(E_{q+1}), O_{q+1})$, and $t_q = \psi(c_q)\psi(c_{q+1})$, for $q = 1, \dots, d$. Then, for $q = 1, \dots, d$,*

$$\alpha(c_q) = \begin{cases} b_{l_q}^{-1} s_{l_q} a_{r_q}, & \textit{if } o(E_q) = 1 \textit{ and } O_q = 1 \\ a_{l_q}^{-1} s_{r_q}^{-1} b_{r_q}, & \textit{if } o(E_q) = 1 \textit{ and } O_q = -1 \\ a_{l_q}^{-1} a_{r_q}, & \textit{if } o(E_q) = -1 \textit{ and } O_q = 1 \\ b_{l_q}^{-1} s_{l_q} s_{r_q}^{-1} b_{r_q}, & \textit{if } o(E_q) = -1 \textit{ and } O_q = -1 \end{cases},$$

$$V_q = (\varepsilon, \delta : \gamma, \delta\gamma), \textit{ for some } \varepsilon, \delta, \gamma = \pm 1,$$

and

(i) $t_q = b_{l_q}^{-1} s_{l_q} s_{l_{q+1}} a_{r_{q+1}}$, *if* $V_q = (1, 1; 1, 1)$,

(ii) $t_q = a_{l_q}^{-1} s_{r_q}^{-1} s_{r_{q+1}}^{-1} b_{r_{q+1}}$, *if* $V_q = (1, -1; 1, -1)$,

(iii) $t_q = b_{l_q}^{-1} s_{l_q} s_{l_{q+1}} s_{r_{q+1}}^{-1} b_{r_{q+1}}$, *if* $V_q = (1, 1; -1, -1)$,

(iv) $t_q = a_{l_q}^{-1} s_{r_q}^{-1} a_{r_{q+1}}$, *if* $V_q = (1, -1; -1, 1)$,

(v) $t_q = a^{-1}_{l_q} s_{l_{q+1}} a_{r_{q+1}}$, *if* $V_q = (-1, 1; 1, 1)$,

(vi) $t_q = b^{-1}_{l_q} s_{l_q} s^{-1}_{r_q} s^{-1}_{r_{q+1}} b_{r_{q+1}}$, *if* $V_q = (-1, -1; 1, -1)$,

(vii) $t_q = a^{-1}_{l_q} s_{l_{q+1}} s^{-1}_{r_{q+1}} b_{r_{q+1}}$, *if* $V_q = (-1, 1; -1, -1)$ *and*

(viii) $t_q = b^{-1}_{l_q} s_{l_q} s^{-1}_{r_q} a_{r_{q+1}}$, *if* $V_q = (-1, -1; -1, -1)$.

Proof. If $o(E_q) = 1$ then

$$\psi(c_q) = z_{\mu_q}(E_q)^{O_q} = \left[b^{-1}_{\nu_q} s_{\nu_q} a_{\nu_q}\right]^{O_q},$$

and the expressions of $\alpha(c_q)$ when $o(E_q) = 1$, follow. When $o(E_q) = -1$ then

$$\psi(c_q) = z_{\mu(O_q)}(E_q)^{O_q \nu^{-1}(l_q)} = \begin{cases} [a_2(E_q)^{-1} a_1(E_q)]^{\nu^{-1}(l_q)}, & \text{if } O_q = 1, \\ [b_1(E_q)^{-1} s_1(E_q) s_2(E_q)^{-1} b_1(E_q)]^{-\nu^{-1}(l_q)}, & \text{if } O_q = -1, \end{cases}$$

and the given expressions for $\psi(c_q)$ when $o(E_q) = -1$ follow, on considering the two possibilities, $l_q = 1$ and $l_q = 2$. That V_q is determined by its first three entries, as claimed, follows from the definition of O_q.

To compute t_q, observe that, from Lemma 2.15, if $O_q = 1$ then U' contains $E^{\varepsilon_q}_{q,l_q} E^{-\varepsilon_{q+1}}_{q+1,l_{q+1}}$, so that $a_{r_q} = b_{l_{q+1}}$. Similarly, if $O_1 = -1$ then U' contains $E^{\varepsilon_{q+1}}_{q+1,l_{q+1}} E^{-\varepsilon_q}_{q,l_q}$, so $a_{l_{q+1}} = b_{r_q}$. To see that (i) holds note that when $o(E_q) = o(E_{q+1}) = O_q = O_{q+1} = 1$, we have $l_q = \nu_q$, $r_q = \mu_q$, $l_{q+1} = \nu_{q+1}$ and $r_{q+1} = \mu_{q+1}$. As $O_q = 1$ we have $a_{r_q} = b_{l_{q+1}}$, so

$$t_q = [b_{\nu_q} s_{\nu_q} a_{\mu_q}][b_{\nu_{q+1}} s_{\nu_{q+1}} a_{\mu_{q+1}}] = [b^{-1}_{l_q} s_{l_q} a_{r_q}][b^{-1}_{l_{q+1}} s_{l_{q+1}} a_{r_{q+1}}] = b^{-1}_{l_q} s_{l_q} s_{l_{q+1}} a_{r_{q+1}}.$$

The other seven cases follow similarly. □

Lemma 4.14. *Let v be an internal vertex of Γ_S and (as on page 44) let $B'_i = \{v^i_1, \dots, v^i_t\}$ be the set of vertices of Γ_S corresponding to the equivalence class B_i, and let g_i be the S-genus of B_i. Let C_v and C^i_j be the cycles extending v and v^i_j, respectively, in Γ''_U. Then the pair (Γ''_U, ψ) is*

1. *a genus 0 joint extension on v by the cyclic word $\psi(C_v)$ and*
2. *a genus g_i joint extension on B'_i by the words $\psi(C^i_1), \dots, \psi(C^i_{t_i})$.*

Proof. First let u be any vertex of Γ_U, with incidence sequence $O_1, \dots, O_d$ and let $E^{\varepsilon_1}_1, \dots, E^{\varepsilon_d}_d$ be the vertex sequence of $\mathrm{lk}(u)$. Let $C_u = c_1 \cdots c_d$ be the cycle extending u. It follows from Lemma 4.13 that all the words a_{l_q}, a_{r_q}, b_{l_q} and b_{r_q} cancel in the product $\psi(C_u) = \psi(c_1) \cdots \psi(c_d)$. Thus

$$\psi(C_u) = \hat{\psi}(c_1) \cdots \hat{\psi}(c_d),$$

where

$$\hat{\psi}(c_q) = \begin{cases} s_{l_q}, & \text{if } o(E_q) = 1 \text{ and } O_q = 1 \\ s_{r_q}^{-1}, & \text{if } o(E_q) = 1 \text{ and } O_q = -1 \\ 1, & \text{if } o(E_q) = -1 \text{ and } O_q = 1 \\ s_{l_q} s_{r_q}^{-1}, & \text{if } o(E_q) = -1 \text{ and } O_q = -1 \end{cases}.$$

If u an internal vertex of Γ_S then, by definition, $s_{l_q} = s_{r_q} = 1$, for all q. Therefore, for the internal vertex v of Γ_S we have $\psi(C_v) = 1$. Moreover, if v is an internal vertex of Γ_S then it is also a vertex of Γ_W and hence has degree at least 3, unless $U = A^{\pm 2}$, for some $A \in \mathcal{A}$. Thus (i) and (ii) of Definition 2.17 hold, and 1 follows.

Assume now that $u = v^i_j \in B'_i$. If $o(E_q) = \varepsilon$, for fixed $\varepsilon = \pm 1$, for all q, then it follows from Lemma 4.12 that $\psi(C_u)$ is conjugate in H to $\theta(W^i_j)$. Hence we may assume that $o(E_q) = 1$ for at least one index q and that $o(E_q) = -1$, for at least two indices q. In this case we may, by renumbering the E_q if necessary, choose E_1 such that $o(E_1) = -1$ and $O_1 = 1$. Let $1 = i_1 < i_2 < \cdots < i_{2m} \le d$ be integers such that $o(E_q) = -1$ if and only if $q \in \{i_1, \ldots, i_{2m}\}$. Then $O_q = -1$ if $i_{2j} \le q < i_{2j+1}$, and $O_q = 1$ if $i_{2j+1} \le q < i_{2(j+1)}$, for $j = 1, \ldots, m$ (subscripts modulo $2m$). Therefore

$$\begin{gathered} \hat{\psi}(c_{i_1}) = \hat{\psi}(c_{i_3}) = \cdots = \hat{\psi}(c_{i_{2m-1}}) = 1, \\ \hat{\psi}(c_{i_{2j}}) = s_{l_{i_{2j}}} s^{-1}_{r_{i_{2j}}}, \text{ for } j = 1, \ldots, m, \\ \hat{\psi}(c_q) = s_{l_q}, \text{ for } i_{2j+1} < q < i_{2(j+1)}, \text{ and} \\ \hat{\psi}(c_q) = s^{-1}_{r_q}, \text{ for } i_{2j} < q < i_{2j+1}. \end{gathered}$$

Hence, for $j = 1, \ldots, m$,

$$\hat{\psi}(c_{i_{2j-1}}) \cdots \hat{\psi}(c_{i_{2j+1}-1}) = s_{l_{i_{2j-1}+1}} \cdots s_{l_{i_{2j}}} s^{-1}_{r_{i_{2j}}} \cdots s^{-1}_{r_{i_{2j+1}-1}},$$

where the subscripts of i are integers modulo $2m$. From Lemma 4.12 again, it follows that $\psi(C_{v^i_j})$ is conjugate in H to $\theta(W^i_j)$.

This is true for all vertices of B'_i, so if $B'_i = \{v^i_1, \ldots, v^i_{t_i}\}$ then

$$\text{genus}_H(\psi(C^i_1), \ldots, \psi(C^i_{t_i})) = \text{genus}_H(\theta(W^i_1), \ldots, \theta(W^i_{t_i})) = g_i - t_i + 1,$$

where g_i is the S-genus of B_i. Thus (i) of Definition 2.17 holds for the extension of B'_i by $\psi(C^i_1), \ldots, \psi(C^i_{t_i})$.

To verify (ii), suppose that $g_i = 0$, $t_i = 1$ and that the degree of v^i_1 is d. Then $\text{genus}_H(\psi(C^i_1)) = \text{genus}_H(\theta(W^i_1)) = 0$ and Lemma 4.10 implies that $\text{genus}_{F(\mathcal{A})}(W^i_1) = 0$. If $d = 1$ then $W^i_1 = 1$. But $W^i_1 = S(E_1^{\varepsilon_1})$ is a cyclic subword of the Wicks form W (or it's inverse) and by definition W is cyclically reduced. Therefore this can't occur.

Now suppose that $d = 2$. Then v^i_j has link $E_1^{\varepsilon_1}, E_2^{\varepsilon_2}$, and since there are an even number of non-alternating edges incident to every vertex of Γ_W, $o(E_1) = o(E_2) = \pm 1$. Moreover W contains $(E_1^{\varepsilon_1} S_1'^{O_1} E_2^{-\varepsilon_2})^{O_1}$ and $(E_2^{\varepsilon_2} S_2'^{O_2} E_1^{-\varepsilon_1})^{O_2}$, where S'_q is defined in the proof of Lemma 4.12. If $o(E_1) = o(E_2) = 1$, then $O_1 = O_2$ and $W^{\pm 1}$ contains $E_1^{\varepsilon_1} S'_1 E_2^{-\varepsilon_2}$ and $E_2^{\varepsilon_2} S'_2 E_1^{-\varepsilon_1}$. In this case the label of the cyclic word W^i is $(S_1'S_2')^{\pm 1}$, so $S_1' = S_2'^{-1}$. Hence W is redundant, a contradiction. In the case $o(E_1) = o(E_2) = -1$, we may assume that $O_1 = -O_2 = 1$ and so $W^{\pm 1}$ contains $E_1^{\varepsilon_1}S_1'E_2^{-\varepsilon_2}$ and $E_1^{\varepsilon_1}S_2'E_2^{-\varepsilon_2}$. In this case the boundary label of W^i is $(S_1'S_2'^{-1})^{\pm 1}$, so W contains two occurrences of $E_1^{\varepsilon_1}S_1'E_2^{-\varepsilon_2}$, which again implies redundancy.

Therefore part (ii) of Definition 2.17 holds. Thus, we have the required genus g_i extension of B_i'. □

We are now in a position to complete the main part of the proof of Theorems 2.23 and 2.24, by showing that (Γ_U'', ψ) is the required extension of U over H. Once this has been done the remaining step is to bound the length of R, and this is addressed in Section 4.4.

Proposition 4.15. *Let* $\text{genus}(S) = k$ *and let* $g = n - k$*. Then the pair* (Γ_U'', ψ) *is a genus* g *extension of* U *over* H*, of length at most* $2K(n)(12l + M + 4)$.

Proof. Let $C_1, \ldots, C_Q$ be the cycles extending the vertices of Γ_U corresponding to the Q boundary components of S; and let $C_{Q+1}, \ldots, C_{Q+t}$ be the cycles extending the internal vertices of S. Then the length of (Γ_U'', ψ) is

$$\sum_{i=1}^{Q+t} |\psi(C_i)|.$$

For $1 \leq j \leq Q + t$, we have $\psi(C_j)$ equal to a product of terms of the form $a_j(A)$, $b_j(A)$ and $s_j(A)$, where A is a long edge of W, and $j = 1$ or 2. From the proof of Lemma 4.11, in particular equations (4.16), (4.17), (4.18) and (4.19), for each long edge A of W and for $j = 1, 2$, each of $a_j(A)$, $b_j(A)$ and $s_j(A)$ occurs in exactly one $\psi(c)$, for some letter c of one of the cycles C_v.

Every $s_j(A)$ is the product of the labels of the short edges between two consecutive long edges of W; and short edges have labels in $F(X)$ of length at most $12l + M + 4$. Each $a_j(A)$ and $b_j(A)$ is a word of length at most $5l + M + 3$ arising from a long edge of W (Lemma 4.5).

By Lemma 2.10, the maximum number of letters in W is $K = \max\{2, 12n - 6\}$. Thus the sum of number of short letters and long letters is at most K. Therefore, if $\mathcal{S}$ and $\mathcal{L}$ are the numbers of short edges and long edges of W, respectively, the length of the extension is bounded by

$$(12l + M + 4)\mathcal{S} + 2(5l + M + 3)\mathcal{L} \quad \leq \quad 2K(12l + M + 4).$$

From Lemma 4.14 (Γ_U'', ψ) is an extension of genus $\sum_{i=1}^{p} g_i$, and the proposition now follows from Lemma 4.10. □

Combining equation (4.11) and Proposition 4.15, we have the required extension and it remains only to verify the bound on the length of R.

4.4 Bounding the length of R

Now assume $h = RFR^{-1}$, where h satisfies the hypotheses and F the conclusions of Theorem 2.23 or Theorem 2.24. We shall consider the geodesic quadrilateral with sides labelled F, R^{-1}, h^{-1}, R, based at 1 in $\Gamma_X(H)$, and refer to these geodesics by their labels. We apply Lemma 2.3 to this quadrilateral, with $\gamma_0 = F$. This gives a partition $F_\iota, F_\gamma, F_\tau$ of F such that F_τ^{-1} δ-fellow travels with an initial segment of R^{-1}, F_ι^{-1} δ-fellow travels with a terminal segment of R and F_γ^{-1}, which may be empty, 2δ-fellow travels with a segment of h, as in Figure 4.7a. Assume first that F satisfies 1 of Theorem 2.23 or Theorem 2.24. If $|F| \le 12l + M + 4\delta + 5$ then, from Lemma 2.2, there is R such that $h =_H RFR^{-1}$ and $|R| \le |h|/2 + 6l + 3M/2 + 2\delta + 7/2$, as claimed. Therefore we assume that $|F| > 12l + M + 4\delta + 5$ and that $F = \theta(W)$, where W is a Wicks form of the appropriate type, with $|\theta(E)| \le 12l + M + 4$, for all $E \in \mathrm{supp}(W)$. Moreover, we assume that R is minimal, in the sense that, for all $(W', \theta') \in \mathcal{F}$, with no long edges, if $h =_H R'\theta'(W')R'^{-1}$ then $|R| \le |R'|$.



(a) A partition $F_\iota, F_\gamma, F_\tau$ (b) p lies on F_τ and q on R^{-1} (c) p lies on F_γ and q on h

Figure 4.7

Let p be a vertex on F such that $|d(p, \iota(F) - d(p, \tau(F)| \le 1$. First suppose that p lies on F_ι or F_τ, so there is a vertex q on R or R^{-1} such that $d(p, q) \le \delta$. Without loss of generality, we may take q to lie on R^{-1} and we take d to be the label of a geodesic from q to p. Then $R = R_0R_1$, where $\tau(R_0) = \iota(R_1) = q$ and $F = F_0F_1$, where $\tau(F_0) = \iota(F_1) = p$, as in Figure 4.7b. This means $|R_1| \ge |F_1| - \delta$ and as $|F_1| \ge (|F| - 1)/2$, we have $|R_1| > 6l + M/2 + \delta + 2$. However $|\theta(E)| \le 12l + M + 4$, for all $E \in \mathrm{supp}(W)$, so p is distance at most $6l + M/2 + 2$ from the initial vertex of $\theta(E^{\pm 1})$, for some $E \in \mathrm{supp}(W)$. Thus $a^{-1}F_1F_0a = \theta(W')$, for some cyclic permutation W' of W and some $a \in F(X)$ with $|a| \le 6l + M/2 + 2$ (see Figure 4.7b). Let $F' = \theta(W')$. As $R_1 = dF_1$, we have $h =_H R_0dF_1F_0d^{-1}R_0^{-1} = R_0daF'a^{-1}d^{-1}R_0^{-1}$ and $|R_0da| \le |R_0| + |a| + |d| \le |R_0| + 6l + M/2 + \delta + 2 < |R_0| + |R_1| = |R|$. As W' is a Wicks form we may replace (W, θ) and F, in the conclusion of Theorem 2.23 or Theorem 2.24, by (W', θ) and F' and then replace R by R_0da. However, this contradicts the minimality of $|R|$. Therefore we may assume that p lies on F_γ.

As $p \in F_\gamma$ there is a vertex q on h such that $d(p, q) \le 2\delta$, and we now take d to be the label of a geodesic from such a vertex q on h to the vertex p on F, as in Figure 4.7c. Let

$h = h_0 h_1$, where $\tau(h_0) = q = \iota(h_1)$; so $h = h_2 h_1 h_0 h_2^{-1}$, for some h_2 such that $|h_2| \leq |h|/2$ (i.e. $h_2 = h_0$ or h_1^{-1}). Then $h_1 h_0 = dF_1 F_0 d^{-1}$ so $h = h_2 dF_1 F_0 d^{-1} h_2^{-1} = h_2 daF' a^{-1} d^{-1} h_2^{-1}$, with a and F' as before. Now $|h_2 da| \leq |h|/2 + 2\delta + 6l + M/2 + 2$ and by hypothesis $|R| \leq |h_2 da|$, so in all cases we have $|R| \leq |h|/2 + 6l + 3M/2 + 7/2 + 2\delta$.

Now assume that F satisfies 2 of Theorem 2.23 or Theorem 2.24. If R is an H-minimal element of $F(X)$ such that $h =_H RFR^{-1}$, where $F = \theta(W)$, then we say that R is a *conjugator* for (W, θ). Since we assumed that W ends in a long edge we call R a *minimal conjugator* if R is a conjugator for (W, θ) and, for every cyclic permutation W' of W which ends in a long edge, and every conjugator R' of (W', θ), we have $|R| \leq |R'|$. Assume then that R is a minimal conjugator. Then F has a partition $\alpha_1, \ldots, \alpha_f$, such that 1 and 2 of Lemma 4.5 hold (with α_i in place of F_i). Let $P = \{\tau(\alpha_i) : 1 \leq i \leq f - 1\}$. Then, from Lemma 4.5 again, $P \neq \emptyset$. Suppose that p lies on F_ι or F_τ, for all $p \in P$. Then, without loss of generality we may assume that p lies on F_τ, for some $p \in P$; so there exists a point q on R^{-1} such that $d(p, q) \leq \delta$. Factorise $R = R_0 R_1$ and $F = F_0 F_1$, as in the previous case. This time $F_1 F_0$ is the label of a Hamiltonian cycle on a graph corresponding to the extension given by the theorem, and the point p is, by definition $v(U_i)$ for some long edge U_i. Thus $F_1 F_0$ is the Hamiltonian cycle of an extension constructed from a permutation W' of W ending with a long edge. Hence we may replace F by $F_1 F_0$ in the conclusion of the theorem. We have $|F_1| \geq 6l + 2$, since α_f is a subinterval of F_1, so $|R_1| \geq 6l + 2 - \delta$, and $h =_H R_0 dF_1 F_0 d^{-1} R_0$, where $|d| \leq \delta < |R_1|$, contrary to the choice of $|R|$ as a minimal conjugator. This contradiction shows that p lies on F_γ, for all $p \in P$.

Now choose $p \in P$, so the geodesic d from some point q on h^{-1} to p has length at most 2δ. Now d is a conjugator, for a cyclic permutation of W ending in a long edge, and by minimality of R we have $|R| \leq |h|/2 + 2\delta$, as required. □

References

[1] J. M. Alonso, T. Brady, D. Cooper, V. Ferlini, M. Lustig, M. Mihalik, M. Shapiro, and H. Short. Notes on word hyperbolic groups. In *Group theory from a geometrical viewpoint (Trieste, 1990)*, pages 3–63. World Sci. Publ., River Edge, NJ, 1991. Edited by Short.

[2] Roland Bacher and Alina Vdovina. Counting 1-vertex triangulations of oriented surfaces. *Discrete Math.*, 246(1-3):13–27, 2002. Formal power series and algebraic combinatorics (Barcelona, 1999).

[3] Martin R. Bridson and James Howie. Conjugacy of finite subsets in hyperbolic groups. *IJAC*, 15(4):725–756, 2005.

[4] M.R. Bridson and A. Häfliger. *Metric Spaces of Non-Positive Curvature.* Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen. Springer, 1999.

[5] Leo P. Comerford and Charles C. Edmunds. Quadratic equations over free groups and free products. *J. Algebra*, pages 276–297, 1981.

[6] Leo P. Comerford, Jr. and Charles C. Edmunds. Solutions of equations in free groups. In *Group theory (Singapore, 1987)*, pages 347–356. de Gruyter, Berlin, 1989.

[7] M. Culler. Using surfaces to solve equations in free groups. *Topology*, pages 237–300, 1981.

[8] Marc Edward Culler. *Genus of elements in a free group*. ProQuest LLC, Ann Arbor, MI, 1978. Thesis (Ph.D.)–University of California, Berkeley.

[9] Franois Dahmani and Vincent Guirardel. Foliations for solving equations in groups: free, virtually free, and hyperbolic groups. *J Topology*, 3(2):343–404, 2010.

[10] Charles C. Edmunds. On the endomorphism problem for free groups. *Comm. Algebra*, 3:1–20, 1975.

[11] Charles C. Edmunds. On the endomorphism problem for free groups. II. *Proc. London Math. Soc. (3)*, 38(1):153–168, 1979.

[12] É. Ghys and P. de La Harpe. *Sur les groupes hyperboliques d'après Mikhael Gromov:*. Progress in mathematics. Birkhäuser, 1990.

[13] R.Z. Goldstein and E.C. Turner. Applications of topological graph theory to group theory. *Math. Z.*, 165:1–10, 1979.

[14] R. I. Grigorchuk and I. G. Lysionok. A description of solutions of quadratic equations in hyperbolic groups. *Internat. J. Algebra Comput.*, 2(3):237–274, 1992.

[15] R.I. Grigorchuk and P.F. Kurchanov. On quadratic equations in free groups. In Kostrikin A.I Bokut' L.A., Ershov Yu L., editor, *Proceedings of the International Conference on Algebra Dedicated to the Memory of A.I. Mal'cev*, volume 1 of *Contemporary Mathematics - American Mathematical Society*, pages 159–171. American Mathematical Society, 1992.

[16] M. Gromov. Hyperbolic groups. In S.M. Gersten, editor, *Essays in group theory*, volume 8 of *MSRI Publications*, pages 75–263. Springer, 1987.

[17] M. Gromov. Asymptotic invariants of infinite groups. In *Geometric group theory, Vol. 2 (Sussex, 1991)*, volume 182 of *London Math. Soc. Lecture Note Ser.*, pages 1–295. Cambridge Univ. Press, Cambridge, 1993.

[18] Leo P Comerford Jr. Quadratic equations over small cancellation groups. *Journal of Algebra*, 69(1):175 – 185, 1981.

[19] O. Kharlampovich, A. Mohajeri, A. Taam, and A. Vdovina. Quadratic Equations in Hyperbolic Groups are NP-complete. *ArXiv e-prints*, June 2013.

[20] O. Kharlampovich and A. Vdovina. Linear estimates for solutions of quadratic equations in free groups. *International Journal of Algebra and Computation*, 22(01):1250004, 2012.

[21] Igor Lysenok and Alexei Myasnikov. A polynomial bound on solutions of quadratic equations in free groups. *Proceedings of the Steklov Institute of Mathematics*, 274(1):136–173, 2011.

[22] I.G. Lysionok. On some algorithmic properties of hyperbolic groups. *Math. USSR Izv.*, 35:145–163, 1990. Translated from Izv. Akad. Nauk. SSSR Ser. Math, vol 54(3), 1989.

[23] G.S. Makanin. Equations in a free group. (russian). *Izv. Akad. Nauk SSSR Ser. Mat.*, 46(6):1199–1273, 1982. English translation in: Math. USSR–Izv. 21 (1983), no. 3, 546–582.

[24] A. Yu. Ol'shanskii. Diagrams of homomorphisms of surface groups. *Sibirsk. Mat. Zh.*, 30(6):150–171, 1989.

[25] Schupp P.E. Quadratic equations in groups, cancellation diagrams on compact surfaces, and automorphisms of surface groups. In Boone W. W. Adian S. I., Higman G., editor, *Word problems II : the Oxford book*. North-Holland Amsterdam, 1979.

[26] A A Razborov. On systems of equations in a free group. *Mathematics of the USSR-Izvestiya*, 25(1):115, 1985.

[27] E. Rips and Z. Sela. Canonical representatives and equations in hyperbolic groups. *Invent. Math.*, 120(3):489–512, 1995.

[28] Alina Vdovina. Products of commutators in free products. *Internat. J. Algebra Comput.*, 7(4):471–485, 1997.

[29] M. J. Wicks. Commutators in free products. *J. London Math. Soc.*, 37:433–444, 1962.

[30] M. J. Wicks. The equation $x^2y^2 = g$ over free products. In *Proceedings of the Second Congress of the Singapore National Acadamy of Science*, pages 238–248, 1973.